

REFERENCES

- Blois, M.S. 1958. *Antioxidant Determination by the Use of a Stable Free Radical*. Nature 181:1199-2000.
- Bushra, S.; Farooq, A.; Muhammad A., 2009. *Effect of Extraction Solvent/Technique on the Antioxidant Activity of Selected Medicinal Plant Extracts*. Molecules, 14: 2167-2180.
- Cook, B. G.; Pengelly, B. C.; Brown, S. D.; Donnelly, J. L.; Eagles, D. A.; Franco, M. A.; Hanson, J.; Mullen, B. F.; Partridge, I. J.; Peters, M.; Schultze-Kraft, R., 2005. *Tropical Forages*. CSIRO, DPI&F(Qld), CIAT and ILRI, Brisbane, Australia. <http://www.tropicalforages.info>, Accessed on November 8, 2013.
- Gomez, S.M., A. Kalamani. 2003. *Butterfly Pea (Clitoria ternatea): A Nutritive Multipurpose Forage Legume for the Tropics - An Overview*. Pakistan Journal of Nutrition 2(6):374-379.
- Grodowska, K., A. Parczewski. 2010. *Organic solvents in the pharmaceutical industry*. Acta Polonaise Pharmaceutics – Drug Research 67(1):3-12.
- Jenshi roobha, J. M. Saravanakumar, K.M. Aravinthan, P. Suganya devi. 2011. *Antioxidant analysis of anthocyanin extracted from Musa acuminata bract*. Journal of Pharmacy Research 4(5):1488-1492.
- Dai, J., R. J. Mumper. 2010. *Plant Phenolics: Extraction, Analysis and Their Antioxidant and Anticancer Properties*. Journal of Molecules 15:7313-7352.
- Luque de Castro, M. D., and L. E. Garcia-Ayuso. 1998. *Soxhlet extraction of solid materials: An outdated technique with a promising innovative future*. Anal. Chimica Acta 369(1-2):1-10.
- Mamidipally, P.K., S.X. Liu. 2004. *First approach on rice bran oil extraction using limonene*. European Journal of Lipid Science and Technology 106(2):122-125.

Masahiro, N., K. Masahiro, N. Minemitsu, K. Akio, N. Yoshimi. 2005. *Scavenging of 1,1-diphenyl-2-picrylhydrazyl (DPPH) by peroxyradical: A Useful Method for Quantitative Analysis of Peroxyradical*. Chem Pharm Bull 53(6):714-716.

Parry, J., L. Su, M. Luther, K. Zhou, M.P. Yurawecz, P. Whittaker, L. Yu. 2005. *Fatty acid composition and antioxidant properties of cold-pressed marionberry, boysenberry, red raspberry, and blueberry seed oils*. J. Agric. Food Chem. 53(3):566-573.

Percival, M. 1998. *'Antioxidants'*, Advanced Nutrition Publications, Inc., Revised.

Peschel W.; Sanchez-Rabaneda F.; Dn W; Plescher A.; Gartzia I.; Jimenez D.; Lamuela-Raventos R.; Buxaderas S.; Condina C. 2006. *An industrial approach in the search of natural antioxidants from vegetable and fruit wastes*. Food Chem., 97: 137-150.

Pinello, M., M. Rubilar, M. Jerez, J. Sineiro, M.J. Nunez. 2005. *Effect of solvent, temperature, and solvent-to-solid ratio on the total phenolic content and antiradical activity of extracts from different components of grape pomace*. Journal of Agricultural Food Chemistry 53(6):2111-2117.

Prakash, A. 2001. *Antioxidant activity*. Medallion Laboratories Analytical Progress 19(2).

Revilla, E., J.M. Ryan, G. Martin-Ortega. 1998. *Comparison of several procedures used for the extraction of anthocyanins from red grapes*. Journal of Agricultural Food Chemistry 46:4592-4597.

Rios, J.L., M.C. Recio. 2005. *Medicinal plants and antimicrobial activity*. Journal of Ethnopharmacology 100(1-2):80-84.

Sanchez-Moreno, C. 2002. *Methods used to evaluate the free radical scavenging activity in foods and biological systems*. Food Sci. Technol Int. 8(3):121-137.

Sasidharan, S., Y. Chen, D. Saravanan, K.M. Sundram, L.Y. Latha. 2011. *Extraction, isolation and characterization of bioactive compounds from plants' extract*.

African Journal of Traditional, Complementary and Alternative Medicines
8(1):1-10.

Spigno, G., L. Tramelli, D.M. Faveri. 2007. *Effects of extraction time, temperature and solvent on concentration and antioxidant activity of grape marc phenolics.* Journal of Food Engineering 81(1):200-208.

Staples, 1992. *Clitoria ternatea L.* Record from Proseabase. Mannerje, L.'t and Jones, R.M. (Editors). PROSEA (Plant Resources of South-East Asia) Foundation, Bogor, Indonesia. http://proseanet.org/prosea/e-prosea_detail.php?frt=&id=17, Accessed on November 8, 2013.

Wang, L., C.L. Weller. 2006. *Recent advances in extraction of nutraceuticals from plants.* Trends in Food Science & Technology 17:300-312.

Yu, L. 2001. *Free radical scavenging properties of conjugated linoleic acids.* J. Agric. Food Chem. 49(7):3452-3456.

Zarnowski, R., Suzuki, Y., Pietr, S.J. 2004. *Alkyl- and alkenylresorcinols of wheat grains and their chemotaxonomic significance.* ZEITSCHRIFT FUR NATURFORSCHUNG C-A Journal of Biosciences 59(3-4) 190-196.

SWISS GERMAN UNIVERSITY