

# Development of Extract Library- IOP

*by* Kholis Audah

---

**Submission date:** 08-Apr-2023 09:13PM (UTC+0700)

**Submission ID:** 2059014205

**File name:** .5\_Audah\_2018\_IOP\_Conf.\_Ser.\_Earth\_Environ.\_Sci.\_130\_012025.pdf (766.45K)

**Word count:** 4308

**Character count:** 24257

**PAPER · OPEN ACCESS****Development of extract library from Indonesian biodiversity: exploration of antibacterial activity of mangrove *bruguiera cylindrica* leaf extracts**

21

To cite this article: K A Audah *et al* 2018 *IOP Conf. Ser.: Earth Environ. Sci.* **130** 012025View the [article online](#) for updates and enhancements.**Related content**

- [Antibacterial activity study of \*Attacus atlas\* cocoon against \*Staphylococcus aureus\* and \*Escherichia coli\* with diffusion and dilution method](#)

Aminah, E R Nugraheni and A Yugutama

26

- [Study on Medicinal Plant Active Substances Extraction and Antibacterial Activity of \*Houttuynia Cordata\*](#)

Ji Yubin, Yang Junjun, Yu Miao *et al.*

24

- [Antibacterial properties of parasitic mistletoe - \*Scurrula ferruginea\* \(Jack\) Danser of Brunei Darussalam](#)

Sheba R David, Amira Amni Adam and Rajan Rajabalaya

13

## Development of extract library from Indonesian biodiversity: exploration of antibacterial activity of mangrove *Bruguiera cylindrica* leaf extracts

K A Audah<sup>1\*</sup>, J Amsyir<sup>1</sup>, F Almasyur<sup>1</sup>, A M Hapsari<sup>1</sup>, and H Sutanto<sup>2</sup>

<sup>1</sup>Department of Biomedical Engineering, Swiss German University, Prominence Tower, West Sutra Road No. 15. Alam Sutra, Tangerang, Banten 15143, Indonesia

<sup>2</sup>Department of Chemical Engineering, Swiss German University, Prominence Tower, West Sutra Road No. 15. Alam Sutra, Tangerang, Banten 15143, Indonesia

\*E-mail: kholis.audah@sgu.ac.id, Tel/ Fax: +622 129779596/ 9598

**Abstract.** Antibacterial drugs derived from natural sources play significant roles in the prevention and treatment of bacterial infections since antibiotics have become less effective against many infectious diseases. Mangroves are very potential natural antibacterial sources among great numbers of wild medicinal plants. *Bruguiera cylindrica* is one of the many mangroves species which spread along Indonesian coastline. The aim of this study was to explore the antibacterial activity of *B. cylindrica* wet and dried leaf extracts. The wet extracts study was conducted with three different solvents system (water, ethanol, and n-Hexane) against *Escherichia coli* and *Staphylococcus aureus*. While, the dried extracts study was conducted with four different solvents system (water, ethanol, chloroform and n-Hexane) against three types of bacteria, *Escherichia coli*, *Staphylococcus epidermidis* and *Staphylococcus aureus*. The study showed that ethanol was the best solvent for extraction of phenolic and flavonoid. Antibacterial activity was measured by zone of inhibition which obtained from agar-disk diffusion method. The widest area of zone of inhibition was showed by wet extracts with ethanol against *S. aureus* and *E. coli* are 14.30 and 13.30 mm, respectively. While, the zone of inhibition dried extracts with ethanol against *S. aureus*, *S. epidermidis* and *E. coli* are 9.32, 6.59 and 6.20 mm, respectively. In conclusion, both type of extracts showed significant antibacterial activity against gram-positive bacteria as crude extracts.

**Keywords:** antibiotics, antimicrobial, infectious diseases, medicinal plants, natural products

### 1. Introduction

Antibiotics are the most important drugs for fighting bacterial infections [1,2]. However, antibiotics have become less effective against many infectious diseases not only because many of them produce toxic reactions to the patient but also due to the emergence of drugs-resistant bacteria [3,4,5,6,7]. The need for new, effective and affordable drugs to treat infectious diseases in the developing world is one of the major issues facing global health today.

Antibacterial drugs derived from natural source play significant roles in the prevention and treatment of human infection diseases [1,8,9]. Natural bioactive compounds would be best solution to overcome the effects of antibiotics resistance bacteria with no or less side effects [10,11]. In recent



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

decades, many studies have been carried out on different plant species to discover compounds of possible interest for medicinal application against bacterial infections [12,13,14]. Among these studies, several have focused on the biological and phytochemical properties of different species of mangrove [15,16,17].

There are about 3.6 million hectares of mangrove forests which grew along 95,000 km of Indonesian coastline and it represent approximately 23% of the total mangrove forests in the world [18,19]. *Bruguiera cylindrica* is one of the many mangroves species which spread along Indonesian coastline. However, the antibacterial potentials of this Indonesian mangroves are yet to be discovered.

Another factor that renewed the exploration of mangrove is the rapid rate of plant species extinction in the past 20 years and the most threatened plant species are found in the tropics [20]. Mangrove forests suffered a shrinkage of about 1-2% every year [21]. Before any medicinal use or other important features can be assessed, it is likely that some species will extinct before they are even discovered.

The aim of this study was to explore the antibacterial activity potency of *B. cylindrica* leaf extracts against *Escherichia coli*, *Staphylococcus aureus* and *Staphylococcus epidermidis*.

## 2. Methods

### 2.1. Plant Materials

The leaves of *B. cylindrica* harvested from Kawasan Ekowisata Mangrove, Pantai Indah Kapuk, North Jakarta, Indonesia with the permission from Department of Marine, Agriculture, and Food Security Special Capital Region of Jakarta. *B. cylindrica* was identified by Indonesian Institute of Sciences (LIPI) taxonomist at Cibinong Science Center. Before used, the leaves were firstly cleaned with tap water to remove dirt and then rinsed with distilled water. The leaves were cut into transverse of approximately 2 cm width and dried by using oven (Memmert, Germany) at 45 °C for 12 hours. The dried leaves were ground into powder using blender (Cucina Philips, Indonesia) and screened by using mesh number 35 (CISA Cedacteria Industrial, Spain). The powder was then transferred into glass bottles with rubber stopper, wrapped in aluminium foil and stored in the freezer at -20 °C prior to extraction.

### 2.2. Extraction

The method was based on method described by Dhayanithi et al. [22] with slightly modification. The leaf powder transferred into an erlenmeyer flask and immersed in solvent (1:10) for 24 hours in a shaker set at 125 rpm. The erlenmeyer flask mouth was sealed with aluminium foil and Parafilm and the whole flask was covered with aluminium foil to protect the content from light. After 24 hours maceration, the extracts were filtrated through cotton fine meshed cloth. The filtrates were then centrifuged (Hettich Rotina 35R centrifuge, Germany) for 10 minutes at 10,000 rpm, 25 °C. The supernatants were then collected by using graduated pipette and prepared for further phytochemical analysis.

### 2.3. Determination of total phenolic content

The determination of total phenolic content (TPC) of extracts obtained was adapted from Banerjee et al., 2008 [23] with slightly modification and calibrated against galic acid as the reference standard. A 0.3 ml sample was mixed with 1.5 ml of Folin-Ciocalteu reagent and 1.2 ml of sodium carbonate (7.5%), consecutively. The mixture of each step should be mixed well by using vortex (Genie 2 mixer, Scientific Industries, USA) and allowed to stand for one hour in a dark chamber. Absorption was measured by using UV-Vis spectrophotometer (Genesys 10 UV-Vis spectrophotometer, Thermo Electron Corporation, USA) at 765 nm. The standard curve galic acid was prepared by diluting the stock standard with the extraction solvents to yield 50 to 200 ppm TPC. The results were calculated according to the calibration curve for galic acid and crude extract of TPC derives from quadruplicate analyses and expressed as galic acid equivalents (GAE mg/ g) of dry material (DM).

#### 2.4. Determination of flavonoid content

The determination of flavonoid content (FC) was adapted from Do et al. [24] with slightly modification and calibrated against quercetin as the reference standard. The Leaf powder was first diluted with its solvent to reach dilution factor of 25 for test the FC in the samples. 1.5 ml of methanol in test tube was prepared. A 0.5 ml sample was mixed with 0.1 ml of 10% aluminium chloride, 0.1 ml of 1 M potassium acetate and 2.8 ml distilled water, consecutively. The mixture of each step should be mixed well by using vortex. The mixtures were incubated in a dark chamber for 30 minutes and absorbance was measured by using a UV-Vis spectrophotometer at 415 nm. The standard curve quercetin was prepared by diluting the stock standard with the extraction solvents to yield 20 - 200 ppm of FC. The results were calculated according to the calibration curve for quercetin and crude extract of FC derives from quadruplicate analyses and expressed as quercetin equivalents (QE mg/ g) of dry material (DM).

#### 2.5. Antibacterial activity

The study of antibacterial activity was conducted in two conditions, the wet extracts study which held at Swiss German University and conducted by using three different solvents (water, ethanol, and n-Hexane) against two types of bacteria, *E. coli* and *S. aureus*. While the dried extracts study which held at Biopharmaca Tropica Research Center, LPPM IPB and conducted by using four different solvents based on the level of polarity (water, ethanol, chloroform and n-Hexane), against three types of bacteria, *E. coli*, *S. epidermidis* and *S. aureus*.

The Agar disk diffusion assay has been widely used to assay plant extract for antibacterial activity [25]. *E. coli*, *S. epidermidis* and *S. aureus* were adjusted to certain concentration,  $1 \times 10^8$  CFU/ml or 0.5 mcFarland's standard [26], were inoculated onto the entire surface of a Mueller-Hinton agar (MHA) plate with a sterile cotton-tipped swab to form an even lawn. In this method, 6 mm sterilized filter papers disks (Whatmann No. 1) were soaked into *B. cylindrica* extract and air dried for one hour at bio safety cabinet. Each type of extracts replicated by three. The paper disks which was impregnated with diluted extracts were placed on the surface of each MHA plate by using a sterile pair of forceps/pincers. Then the plates were incubated at 37°C for 24 hours and diameter of the zone of inhibition (ZOI) obtained was measured by a ruler or caliper. The bigger diameter of the ZOI, the more susceptible of the microorganism to the extracts.

29

#### 2.6. Statistical analysis


Data result was analyzed by Analysis of Variance (ANOVA) single factor. The P-two tail or P-Value below 0.05 ( $P < 0.05$ ) indicated that the data were significantly different, while if P-Value above 0.05 ( $P > 0.05$ ), the data were insignificantly different. The data graphic was built using Microsoft Excel 2010.

### 3. Results and Discussion


Extraction of plant materials depends on various factors such as solvents, methods, and extraction time to separate different quality and quantity of bioactive components in the crude extracts [27]. Polarity of solvent plays key role in the outcome of bioactive compound extracted from *B. cylindrica*. The wet extracts with ethanol showed the highest total phenolic content compared with water and n-Hexane, consecutively (Figure 1). It was because ethanol has the ability to bind with hydrophilic compounds such as flavonoid and phenolic compound [28]. Ethanol is chosen when water-insoluble constituents need to be extracted. However, higher ethanol percentages do not necessarily mean higher extraction activity [29].

The wet extracts with ethanol also showed the highest flavonoid content compared with water and n-hexane, consecutively (Figure 2). Less polar flavonoids (e.g., isoflavones, flavanones, methylated flavones, and flavonols) are best extracted with chloroform, dichloromethane, diethyl ether, or ethyl acetate, while flavonoid glycosides and more polar aglycones are best extracted with alcohols or

alcohol–water mixtures [30,31]. The combination of water and organic solvent facilitate the extraction of chemicals that are soluble in water and/or organic solvent [24].



**Figure 1.** Total phenolic content of *B. cylindrica* leaf extracts




**Figure 2.** Flavonoid content of *B. cylindrica* leaf extracts

The mechanisms for phenolic toxicity to microorganisms include adsorption and disruption of microbial membranes, interaction with enzymes, and metal ion deprivation [32,33,34]. The higher FC was also expected to show the higher antibacterial activity since flavonoids are synthesized by plants in response to microbial infection [35,36,37,38]. Flavonoid involved in inhibition nucleic acid biosynthesis, inactivate microbial adhesins enzymes and cell envelope transport proteins [39,40]. Lipophilic flavonoids may also disrupt microbial membranes [13,36,41].

The ZOI obtained of the wet extracts with ethanol against *S. aureus* and *E. coli* were 14.30 and 13.30 mm, respectively (Figure 3). The dried extracts study also showed a consistent results with the wet extracts study. The ZOI of the dried extracts with ethanol against *S. aureus*, *E. coli*, and *S. epidermidis* were 9.32, 6.20 and 6.59 mm, as showed in Table 1. Taken altogether, the leaf extracts can be categorized as bacteriostatic due to the capability of bacterial growth inhibition. Quantitatively, the leaf extract can be classified as weak-to-moderate antibacterial since the ZOI obtained around 10

to 15 mm. While, the ZOI obtained by Ciprofloxacin, tetracycline and chloramphenicol were 41.2, 22.10 and 20.80 mm.



**Figure 3.** Zone of inhibition of wet extracts

**Table 1.** Antibacterial test of *B. cylindrica* leaf extracts againsts *E. coli* and *S. aureus*

Conc (%)	(+ mm)		Extraction						(-) mm DMSO
	Tetra	Chloram	b. Cyl ethx (mm)			b. Cyl chlx (mm)			
	22.10	20.80	SA	EC	SE	SA	EC	SE	
0.4			9.32	6.20	6.59	7.32	6.88	7.42	
0.2			8.32	6.00	6.32	7.00	6.52	6.26	
0.1			8.08	6.00	6.16	6.72	6.00	6.00	
0.05			6.40	6.00	6.20	6.56	6.02	6.52	
									6.00

Conc : concentration, tetra : tetracycline, chlor : chloramphenicol, b. cyl ethx : *B. cylindrica* ethanol extract, b. cyl chlx : *B. cylindrica* chloroform extract, r. muc ethx : *R. mucronata* ethanol extract, r. muc chlx : *R. mucronata* chloroform extract, DMSO = DMSO 20%, SA: *S. aureus*, EC: *E. coli*, and SE: *S. epidermidis*.

Since all the extracts consist of many different compounds, they presumably contain single compounds with higher activity than the value obtained as crude extract. In addition, isolation and characterization of the antibacterial compounds presented in these extracts seemed to be worthwhile [42]. The potential for developing antimicrobial agents from mangrove species appears rewarding, as it may lead to development of phytomedicine against pathogenic microbes [43].

Study with different sampling time, age of plant, plant part, solvent, method of extraction and time required for extraction would be necessary [44,45]. Because, many of the phytochemicals in plants can be detected at different concentrations because the amount and composition of secondary metabolites are not constant and their concentration depends on the tissue type and the age of the plant [46,47].

In general, both type of extracts showed significant antibacterial activity against gram-positive bacteria. This matter may be correlated with the cell envelopes composition. The cell envelopes of most bacteria fall into one of two major groups. Gram-negative bacteria are surrounded by a thin peptidoglycan cell wall, which itself is surrounded by an outer membrane containing lipopolysaccharide. Gram-positive bacteria lack an outer membrane but are surrounded by layers of peptidoglycan many times thicker than is found in the Gram-negatives [48]. Therefore, gram-negative bacteria is more resistant than gram-positive due to the lipopolyproteins and peptidoglycan

composition in their cell envelope [49]. Gram-negative cell walls is strong to withstand three atm of turgor pressure, tough to endure extreme temperatures, pH and elastic to be capable of expanding several times their normal surface area [50].

### Conclusions

The study of *B. cylindrica* leaf extract suggested that ethanol was the best solvent. The widest area of ZOI was showed by wet extracts with ethanol against *S. aureus* and *E. coli* were 14.30 and 13.30 mm, respectively. While, the ZOI dried extracts with ethanol against *S. aureus*, *S. epidermidis* and *E. coli* were 9.32, 6.59 and 6.20 mm, respectively. In conclusion, both type of extracts showed considerably high antibacterial activity against gram-positive bacteria as crude extracts.

### Acknowledgments

The authors gratefully acknowledge that the present research is supported by the Central Research Fund of the Swiss German University with Contract Number A/REC/0073/IX/2015 and by Ministry of Research and Technology and Higher Education Republic of Indonesia with Contract Number SP DIPA-042.06.1.401516/2017. We also thank Directorate of Marine, Agriculture and Food Security, DKI Jakarta Province for giving us permission to collect mangrove leaves at the Mangroves Sanctuary Region, Pantai Indah Kapuk, North Jakarta, Indonesia. We thank our research collaborators Dr. Irmanida Batubara and staffs at Biopharmaca Tropica Research Center, Bogor Agricultural University and Dr. Doddy Kustaryono and staffs at STKIP Surya, Tangerang. We also thank the Director and staffs of Academic Research and Community Services Swiss German University for their support throughout the project.

### References

- [1] WHO (2014). Antimicrobial resistance: global report on surveillance. WHO Press. Geneva. Switzerland
- [2] CDC (2013). Antibiotic resistance threats in the United States. US Department of Health and Human Services. Atlanta: Georgia
- [3] Kart A, Bilgili A (2008). Ionophore antibiotics: toxicity, mode of action and neurotoxic aspect of carboxylic ionophores. *J Animal Vet Adv.* 7(6):748-751.
- [4] Bassetti M, Ginocchio F, Mikulska M. New treatment options against gram-negative organisms (2011). *Critical Care.* 15: 215
- [5] Grill MF, Maganti RK (2011). Neurotoxic effects associated with antibiotic use: management considerations. *British journal of clinical pharmacology.* 72(3):381-93.
- [6] Drouet M, Chai F, Barthelemy C, Lebuffe G, Debaene B, Decaudin B, et al., (2015). Endothelial Cell Toxicity of Vancomycin Infusion Combined with Other Antibiotics. *Antimicrobial agents and chemotherapy.* 59(8):4901-6.
- [7] Ventola C (2015). The Antibiotic Resistance Crisis Part 1: Causes and Threats. *P & T.* 40(4):277-83.
- [8] Bhalodia NR, Nariya PB, Shukla VJ. (2011). Antibacterial and Antifungal activity from Flower Extracts of *Cassia fistula* L.: An Ethnomedicinal Plant. *International Journal of PharmTech Research.* Vol. 3, No.1, pp 160-168,
- [9] Prihanto AA, Firdaus M, Nurdiani R (2012). Anti-Methicillin Resistant *Staphylococcus aureus* (MRSA) of Methanol Extract of Mangrove Plants Leaf: Preliminary Report. *Drug discovery today.* 4(8):439-440
- [10] Bansal D, Bhasin P, Punia A, Sehrawat AR (2011). Evaluation of antimicrobial activity and phytochemical screening of extracts of *Tinospora cordifolia* against some pathogenic microbes. *J. Pharm. Res.* 5(1):127-129.
- [11] Priyatharsini S, Sivagurunathan P, Uma C, Bhuvanewari M, Aruljothi S (2015). Assessment of antibacterial activity of halophytic plants against uropathogens. *Asian J Pharm Sci Tech.* 5(2): 102-105.



- [12] Chan BCL, Lau CBS, Jolival C, Lui SL, GanemElbaz C, Paris JM, Litaudon M, Fung KP, Leung PC, Ip M. 2011. Chinese medicinal herbs against antibiotic-resistant bacterial pathogens. in Science against microbial pathogens: communicating current research and technological advances. Formatex
- [13] Djuoussi MG, Tamokou JD, Ngnokam D, Kuate JR, Tapondjou LA, Harakat D, et al., (2015). Antimicrobial and antioxidant flavonoids from the leaves of *Oncoba spinosa* Forssk. (Salicaceae). BMC complementary and alternative medicine. 15:134.
- [14] Voukeng IK, Beng VP, Kuete V (2017). Multidrug resistant bacteria are sensitive to *Euphorbia prostrata* and six others Cameroonian medicinal plants extracts. BMC research notes.10(1):321.
- [15] Ravikumar S, Gnanadesigan M, Suganthi P, Ramalakshmi A (2010). Antibacterial potential of chosen mangrove plants against isolated urinary tract infectious bacterial. Int J Med Medical Sci. 2(3): 94-99.
- [16] Saad S, Taher M, Susanti D, Qaralleh H, Awang AFIB (2012). In vitro antimicrobial activity of mangrove plant *Sonneratia alba*. Asian Pacific Journal of Tropical Biomedicine. 2(6):427-9.
- [17] Ramasubburayan R, Prakash S, iyapparaj P, Sumathi S, Thaddaeus BJ, palavesam A, immanuel G (2015). Investigation on antibacterial, antifungal and cytotoxic properties of chosen mangroves. Indian J Geo-Marine Sci. 44(11): 1769-1777.
- [18] Giesen W., Wulffraat S., Zieren M., Scholten L (2006). Mangrove guide book for Southeast Asia. FAO and Wetlands International, Thailand: Dharmasarn Co., Ltd.
- [19] Giri C (2016). Observation and Monitoring of Mangrove Forests Using Remote Sensing: Opportunities and Challenges. Remote Sensing. 8(9):783.
- [20] IUCN 2010. New study shows over one fifth of the world's plants are under threat of extinction. <http://www.iucnredlist.org/news/srli-plants-press-release>
- [21] Locatelli T, Binet T, Kairo JG, King L, Madden S, Patenaude G, et al., (2014). Turning the tide: how blue carbon and payments for ecosystem services (PES) might help save mangrove forests. Ambio. 43(8):981-95.
- [22] Dhayanithi N, Kumar T, Murthy R, Kathiresan K (2012). Isolation of antibacterials from the mangrove, *Avicennia marina* and their activity against multi drug resistant *Staphylococcus aureus*. Asian Pac. J. Trop. Biomed. 2(3):1892-1895.
- [23] Banerjee D, Chakrabarti S, Hazra AK, Banerjee S, Ray J, Mukherjee B (2008). Antioxidant activity and total phenolics of some mangroves in Sundarbans. Afr. J Biotech. 7(6):805-810.
- [24] Do QD, Angkawijaya AE, Tran-Nguyen PL, Huynh LH, Soetaredjo FE, Ismadji S, et al., (2014). Effect of extraction solvent on total phenol content, total flavonoid content, and antioxidant activity of *Limnophila aromatica*. Journal of food and drug analysis. 22(3):296-302.
- [25] Ergene A, Guler P, Tan S, Mirici S, Hamzaoglu E, Duran (2006). Antimicrobial and antifungal activity of *Heracleum sphondylium* subsp. artvinense. Afr J Biotechnol. 5(11):1087-1089.
- [26] Baris O, Gulluce M, Sahin F, Ozer H, Kilic H, Ozkan H, Sokmen M, Ozbek T (2006). Biological activities of the essential oil and methanol extract of *Achillea biebersteinii* Afan. (Asteraceae). Turk. J. Biol. 30:65-73.
- [27] Azmir J, Zaidul ISM, Rahman MM, Sharif KM, Mohamed A, Sahena F, et al., (2013). Techniques for extraction of bioactive compounds from plant materials: A review. Journal of Food Engineering. 117(4):426-36.
- [28] Sasidharan S, Nilawaty R, Xavier R, Latha LY, Amala R (2010). Wound Healing Potential of *Elaeis guineensis* Jacq Leaves in an Infected Albino Rat Model. Molecules. 15(5):3186-99.
- [29] Morgan M (2009). Ethanol in Herbal Medicine. Mediherb 129
- [30] Brglez Mojzer E, Knez Hrnčić M, Skerget M, Knez Z, Bren U (2016). Polyphenols: Extraction Methods, Antioxidative Action, Bioavailability and Anticarcinogenic Effects. Molecules. 21(7).
- [31] Bouhoreira A, Dadamoussa B, Gherraf N (2016). Phytochemical study and evaluation antibacterial activity of flavonoid excerpts: Male spathe of date palm. Der Pharmacia Lettre. 8 (13):171-176

- [32] Fattouch S, Caboni P, Coroneo V, Tuberoso C, Angioni A, Dessi S, Marzouki N, Cabras P (2007). Antimicrobial Activity of Tunisian Quince (*Cydonia oblonga* Miller) Pulp and Peel Polyphenolic Extracts. *J. Agric. Food Chem.* 55(3):963- 969.
- [33] Maddox CE, Laur LM, Tian L (2009). Antibacterial Activity of Phenolic Compounds Against the Phytopathogen *Xylella fastidiosa*. *Current Microbiology.* 60(1):53-8.
- [34] Fu L, Lu W, Zhou X (2016). Phenolic Compounds and In Vitro Antibacterial and Antioxidant Activities of Three Tropic Fruits: Persimmon, Guava, and Sweetsop. *BioMed research international.* 2016:4287461.
- [35] Baba SA, Malik SA (2015). Determination of total phenolic and flavonoid content, antimicrobial and antioxidant activity of a root extract of *Arisaema jacquemontii* Blume. *Journal of Taibah University for Science.* 9(4):449-54.
- [36] Kumar S, Pandey AK (2013). Chemistry and biological activities of flavonoids: an overview. *TheScientificWorldJournal.* 2013:162750.
- [37] Mierziak J, Kostyn K, Kulma A. (2014). Flavonoids as Important Molecules of Plant Interactions with the Environment. *Molecules.* 19(10):16240-65.
- [38] Sofiane G, Wafa N, Abbas K, Amar O (2015). Antioxidant and antimicrobial activities of flavonoids extracted from *Thymus ciliatus* (Desf.) Benth. *Der Pharmacia Lettre.* 7(7):358-363.
- [39] Tkachenko H, Buyun L, Terech-Majewska E, Osadowski Z (2016). Antibacterial activity of ethanolic leaf extracts obtained from various ficus species (Moraceae) against the fish pathogen, *Citrobacter freundii*. *J Ecol Protection of the Coastline.* 20: 117-136.
- [40] Qwarse M, Sempombe J, Mihale MJ, Henry L, Mugoyela V, Sung'hwa F (2017). Cytotoxicity, Antibacterial, and Antifungal Activities of Five Plant Species Used by Agro-pastoral Communities in Mbulu District, Tanzania. *Int J Res Pharm Chem,* 7(1): 1-14.
- [41] Mishra MP, Rath S, Swain SS, Ghosh G, Das D, Padhy RN (2017) In vitro antibacterial activity of crude extracts of 9 selected medicinal plants against UTI causing MDR bacteria. *Journal of King Saud University - Science.* 29(1):84-95.
- [42] Eloff JN, Famakin JO, Katerere DRP (2005). *Combretum woodii* (Combretaceae) leaf extracts have high activity against Gram-negative and Gram-positive bacteria. *Afr. J. Biotechnol.* 4(10):1161-1166.
- [43] Patra JK, Thatoi H, Panigrahi TK, Rath SK, Dhal NK (2009). Phytochemical Screening and Antimicrobial Assessment of Leaf Extracts of *Excoecaria Agallocha* L.: A Mangal Species of Bhitarkanika, Orissa, India. *Adv Natur Appl Sci,* 3(2): 241-246
- [44] Sasidharan S, Chen Y, Saravanan D, Sundram KM, Latha LY (2011). Extraction, isolation and characterization of bioactive compounds from plants extracts. *Afr J Tradit Complement Altern Med.* 8(1):1-10.
- [45] Okoduwa SIR, Umar IA, James DB, Inuwa HM, Habila JD (2016). Evaluation of extraction protocols for anti-diabetic phytochemical substances from medicinal plants. *World Journal of Diabetes.* 7(20):605.
- [46] Silva-Beltrán NP, Ruiz-Cruz S, Cira-Chávez LA, Estrada-Alvarado MI, Ornelas-Paz JdJ, López-Mata MA, et al., (2015). Total Phenolic, Flavonoid, Tomatine, and Tomatidine Contents and Antioxidant and Antimicrobial Activities of Extracts of Tomato Plant. *International Journal of Analytical Chemistry.* 2015:1-10.
- [47] Wouters FC, Blanchette B, Gershenzon J, Vassão DG (2016). Plant defense and herbivore counter-defense: benzoxazinoids and insect herbivores. *Phytochemistry Reviews.* 15(6):1127-51.
- [48] Silhavy TJ, Kahne D, Walker S. The bacterial cell envelope. (2010). *Cold Spring Harbor perspectives in biology.* 2(5):a000414.
- [49] Lopez D, Kolter R. Functional microdomains in bacterial membranes (2010). *Genes & development.* 24(17):1893-902.
- [50] Beveridge TJ. Structures of Gram-Negative Cell Walls and Their Derived Membrane Vesicles (1999). *J Bacteriology.* 18(16): 4725±4733

# Development of Extract Library-IOP

---

## ORIGINALITY REPORT

---

24%

SIMILARITY INDEX

19%

INTERNET SOURCES

17%

PUBLICATIONS

11%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

1	<a href="http://digitalcommons.lsu.edu">digitalcommons.lsu.edu</a> Internet Source	1%
2	<a href="http://shodh.inflibnet.ac.in:8080">shodh.inflibnet.ac.in:8080</a> Internet Source	1%
3	<a href="http://www.bcc.bas.bg">www.bcc.bas.bg</a> Internet Source	1%
4	Submitted to American Intercontinental University Online Student Paper	1%
5	Submitted to University of Nottingham Student Paper	1%
6	<a href="http://jfas.info">jfas.info</a> Internet Source	1%
7	<a href="http://www.aensiweb.com">www.aensiweb.com</a> Internet Source	1%
8	Submitted to MAHSA University Student Paper	1%
9	<a href="http://www.excli.de">www.excli.de</a> Internet Source	1%

---

10	Lim, Y.Y.. "Antioxidant properties of Phyllanthus amarus extracts as affected by different drying methods", LWT - Food Science and Technology, 200711 Publication	1 %
11	jels.ub.ac.id Internet Source	1 %
12	link.springer.com Internet Source	1 %
13	repository.sgu.ac.id Internet Source	1 %
14	Shoib A. Baba, Shahid A. Malik. " Determination of total phenolic and flavonoid content, antimicrobial and antioxidant activity of a root extract of Blume ", Journal of Taibah University for Science, 2018 Publication	1 %
15	prosiding.farmasi.unmul.ac.id Internet Source	1 %
16	Submitted to Our Lady of Fatima University Student Paper	<1 %
17	cmuj.cmu.ac.th Internet Source	<1 %
18	dspace.kuet.ac.bd Internet Source	<1 %

19	<a href="http://www.academicjournals.org">www.academicjournals.org</a> Internet Source	<1 %
20	Submitted to University of Newcastle Student Paper	<1 %
21	<a href="http://docobook.com">docobook.com</a> Internet Source	<1 %
22	<a href="http://docslib.org">docslib.org</a> Internet Source	<1 %
23	<a href="http://journals.univ-tlemcen.dz">journals.univ-tlemcen.dz</a> Internet Source	<1 %
24	<a href="http://www.imedpub.com">www.imedpub.com</a> Internet Source	<1 %
25	<a href="http://www.researchgate.net">www.researchgate.net</a> Internet Source	<1 %
26	<a href="http://www.science.gov">www.science.gov</a> Internet Source	<1 %
27	<a href="http://bm Cresnotes.biomedcentral.com">bm Cresnotes.biomedcentral.com</a> Internet Source	<1 %
28	Mangrove Ecosystems of Asia, 2014. Publication	<1 %
29	<a href="http://iosrjournals.org">iosrjournals.org</a> Internet Source	<1 %
30	<a href="http://nibio.brage.unit.no">nibio.brage.unit.no</a> Internet Source	<1 %

31

[www.intechopen.com](http://www.intechopen.com)

Internet Source

&lt;1 %

32

"Nanotechnology: Applications in Energy, Drug and Food", Springer Science and Business Media LLC, 2019

Publication

&lt;1 %

33

I.M.C. Brighente, M. Dias, L.G. Verdi, M.G. Pizzolatti. "Antioxidant Activity and Total Phenolic Content of Some Brazilian Species", Pharmaceutical Biology, 2008

Publication

&lt;1 %

34

Traba, Christian, Long Chen, and Jun F. Liang. "Low power gas discharge plasma mediated inactivation and removal of biofilms formed on biomaterials", Current Applied Physics, 2013.

Publication

&lt;1 %

35

[discovery.ucl.ac.uk](http://discovery.ucl.ac.uk)

Internet Source

&lt;1 %

36

[expert.ubd.edu.bn](http://expert.ubd.edu.bn)

Internet Source

&lt;1 %

37

[msptm.org](http://msptm.org)

Internet Source

&lt;1 %

38

[nopr.niscpr.res.in](http://nopr.niscpr.res.in)

Internet Source

&lt;1 %

39

[research-repository.griffith.edu.au](http://research-repository.griffith.edu.au)

<1 %

40

[ujcontent.uj.ac.za](http://ujcontent.uj.ac.za)

Internet Source

<1 %

41

[works.bepress.com](http://works.bepress.com)

Internet Source

<1 %

42

[www.revistabionatura.com](http://www.revistabionatura.com)

Internet Source

<1 %

43

Dhayanithi, Nagarajan Balachandran, Thipramalai Thankappan Ajith Kumar, Jesu Arockiaraj, Chellam Balasundaram, and Ramasamy Harikrishnan. "Dietary supplementation of *Avicennia marina* extract on immune protection and disease resistance in *Amphiprion sebae* against *Vibrio alginolyticus*", *Fish & Shellfish Immunology*, 2015.

Publication

<1 %

44

Muhammad Luqman Nordin, Arifah Abdul Kadir, Zainul Amiruddin Zakaria, Rasedee Abdullah, Muhammad Nazrul Hakim Abdullah. "In vitro investigation of cytotoxic and antioxidative activities of *Ardisia crispa* against breast cancer cell lines, MCF-7 and MDA-MB-231", *BMC Complementary and Alternative Medicine*, 2018

Publication

<1 %

45 Prachi Jha, Ashish Saraf, Jasmeet Kaur Sohal. <1 %  
"Antimicrobial Activity of Biologically  
Synthesized Gold Nanoparticles from Wild  
Mushroom Cantharellus Species", Journal of  
Scientific Research, 2021  
Publication

---

46 Vijayarathna, Soundararajan, Zuraini Zakaria,  
Yeng Chen, Lachimanan Yoga Latha, Jagat R.  
Kanwar, and Sreenivasan Sasidharan. "The  
Antimicrobial Efficacy of Elaeis guineensis:  
Characterization, in Vitro and in Vivo Studies",  
Molecules, 2012.  
Publication

---

47 doaj.org <1 %  
Internet Source

---

48 journals.jkuat.ac.ke <1 %  
Internet Source

---

49 jurcon.ums.edu.my <1 %  
Internet Source

---

50 Aditya U. Arvindekar, Galvina R. Pereira, Kirti  
S. Laddha. "Assessment of conventional and  
novel extraction techniques on extraction  
efficiency of five anthraquinones from Rheum  
emodi", Journal of Food Science and  
Technology, 2015  
Publication

---



51

Antonio C. Santos Felix, Cleber G. Novaes, Maísla Pires Rocha, George E. Barreto et al. "Mixture Design and Doehlert Matrix for the Optimization of the Extraction of Phenolic Compounds from Spondias mombin L Apple Bagasse Agroindustrial Residues", *Frontiers in Chemistry*, 2018

Publication

&lt;1 %

52

J.N. Eloff, D.R. Katerere, L.J. McGaw. "The biological activity and chemistry of the southern African Combretaceae", *Journal of Ethnopharmacology*, 2008

Publication

&lt;1 %

53

P. Gopalakrishnakone. "Therapeutic Potential of Plants as Anti-microbials for Drug Discovery", *Evidence-based Complementary and Alternative Medicine*, 06/24/2008

Publication

&lt;1 %

54

Maria Fernanda Taviano, Khaled Rashed, Angela Filocamo, Francesco Cacciola et al. "Phenolic profile and biological properties of the leaves of *Ficus vasta* Forssk. (Moraceae) growing in Egypt", *BMC Complementary and Alternative Medicine*, 2018

Publication

&lt;1 %

55

Sami Fattouch, Pierluigi Caboni, Valentina Coroneo, Carlo I. G. Tuberoso et al. "Antimicrobial Activity of Tunisian Quince (

&lt;1 %

Miller) Pulp and Peel Polyphenolic Extracts ",  
Journal of Agricultural and Food Chemistry,  
2007

Publication

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On