

# Iconiet-2018

*by* Kholis Audah

---

**Submission date:** 05-Jan-2021 09:57AM (UTC+0700)

**Submission ID:** 1483145565

**File name:** ICONIET\_2018\_-\_Kholis\_Abdurachim\_Audah\_Full\_Paper\_1.docx (351.47K)

**Word count:** 3489

**Character count:** 18619

# Anticancer Screening of Mangrove Extract Library: Accelerating Drug Discovery from Indonesian Biodiversity

Kholis Abdurachim AUDAH<sup>a,b\*</sup>, Razethy BATUBARA<sup>a</sup>, Julkipli AMSYIR<sup>a</sup>, Sandiego HIMAWAN<sup>a</sup> and Irmanida BATUBARA<sup>c,d</sup>

<sup>a</sup>Department of Biomedical Engineering, Faculty of Life Sciences and Technology, Swiss German University, Tangerang 15143, Indonesia, <sup>b</sup>Academic Research and Community Services, Swiss German University, Tangerang 15143, Indonesia, <sup>c</sup>Biopharmaca Tropica Research Center, Bogor Agricultural University, Bogor 16128, Indonesia, <sup>d</sup>Department of Chemistry, Faculty of Mathematics and Natural Sciences, Bogor Agricultural University, 16680

\*Corresponding author e-mail: audahka@gmail.com

**Abstract:** Cancer is still one of the leading death causes in the world. There will be an estimated 18.1 million new cancer cases and over 9 million cancer deaths in 2018. Furthermore, some studies reported that new types of cancer had been discovered. Therefore, it is very important to find different sources for anticancer agents. Indonesian biodiversity is abundant natural resources that can be utilized as potential drug sources. Mangroves are among potential plant medicine that grow nearly at all Indonesian coastlines. The aim of this study was to evaluate the potential of mangrove extracts (extract library) as anticancer agents. In this study, eight mangroves species were used. There were 16 samples collected from different parts of the plants such as leaf, bark or root. Four types of solvents with different polarity were used producing 64 extracts. Brine Shrimp Lethality Test (BSLT) and MTT Assay were conducted for anticancer screening of mangrove extract library. There were 43 extracts showed anticancer potential with  $LC_{50} < 1000$  ppm and 21 extracts with  $LC_{50} > 1000$  ppm. Next, 16 out of 43 extracts which have  $LC_{50} < 500$  ppm and some extracts with  $LC_{50} > 1000$  ppm were selected to be evaluated in MTT Cytotoxicity assay. There were 3 extracts that highly inhibited HeLa and MCF-7 Cell line. The highest inhibition percentage was recorded for ethyl acetate extract of root of *Avicennia Marina* (85 Ea), although it was not toxic enough in BSLT due to the high  $LC_{50}$  value ( $>1000$  ppm). The next highest inhibition percentage were recorded for water extract of leaves of *Sonneratia alba* (88 A) and ethyl acetate extract of leaves of *Xylocarpus granatum* (86 Ea). In addition, there were 2 extracts that only effective to inhibit HeLa cell line, namely ethanol extract of leaves *Bruguiera gymnorrhiza* (75 Et) and ethyl acetate extract of root of *Rhizophora apiculata* (74 Ea). Phytochemical analysis of the extracts was also evaluated. The majority of samples showed saponin and tannin in considerable amount. This supported the data that mangrove extracts were potential as anticancer agents.

**Keywords:** Anticancer, drug discovery, drug resistant, extract library, Indonesian biodiversity.

22

## 1. Introduction

Cancer is still one of the leading death causes in the world. There will be an estimated 18.1 million new cancer cases and over 9 million cancer deaths in 2018 (Bray *et al.*, 2018). Lung cancer is the most frequently diagnosed type of cancer and become the leading cause of cancer deaths in male and female. On the other hand, effective cancer therapy is still a great challenge for nowadays society due to some limitations and side-effects from the treatment. Therefore, finding alternatives for drug sources as anticancer agent is urgently required. Meanwhile, current anticancer research shows that herbal plants from tropical region such as mangrove have a potential cytotoxicity effects towards cancer cells (Kerry *et al.*, 2018). In addition, anticancer drug discovery is a lengthy and expensive process. Through screening process utilizing natural products can become a solution of the slow and expensive drug discovery process using conventional way. Indonesia is well known as one of the richest countries in

the world in biodiversity where is a home of approximately 11% or more than 30,000 of the world's flowering plants and other biota both in land and marine with significant figures (Rintelen, Arida, & Häuser, 2017). One of potential plants as medicinal sources and widely spread along Indonesian coastline is Mangrove. Mangrove and mangrove associates are very potential plants as medicinal sources (Bandaranayake, 2002; Audah, 2018). Along roughly 90,000 kilometers coastline, Indonesia is home of about 20 families with hundreds of species of mangroves and their associates. Indonesia has the largest mangrove forest or about 23% of total world mangrove forests (Giri *et al.*, 2011). The aim of this study was to evaluate the potential of mangrove extracts (extracts library) as anticancer agents. Since long time ago different parts of mangrove trees either roots, branches, leaves, flowers and the fruits had been utilized as food and medicinal sources.

2. Research Methods

2.1. Material

The main raw material used in this study was various parts of mangroves plants such as leaves, stem and roots. There were eight different species of mangroves collected from the Eastern Coastline of Lampung Province, Indonesia in May 2018. The species identification was done at the Herbarium of the Center for Biological Research of the Indonesian Institute of Sciences, Bogor, Indonesia. Samples were collected from each parts of the plants in the total of 16 samples.

Table 1: Mangroves species and part of the plants used in this study\*.

Name of Plants	Part of Plants	Samples Code
<i>Rhizophora apiculata</i>	Root	74
<i>Bruguiera gymnorrhiza</i>	Leaf	75
<i>Rhizophora mucronata</i>	Leaf	79
	Stem	80
<i>Avicennia marina</i>	Leaf	84
	Root	85
<i>Xylocarpus granatum</i>	Leaf	86
<i>Ceriops tagal</i>	Leaf	87
<i>Sonneratia caseolaris</i>	Leaf	88

\*Part of plants collected were based upon the nature of the plants.

The chemicals and solvents used in this study were obtained from Brata Chem including: Tween-80, Distilled water, ethanol, n-hexane, ethyl acetate Concentrated HCl, n-amyl alcohol, acetone, dichloromethane, Liebermann-Burchard reagent, chloroform-ammonia, H<sub>2</sub>SO<sub>4</sub> 2 M, Mayer, Dragendorf and Wagner reagents, Magnesium powder, FeCl<sub>3</sub> 1%, NaOH 10% and DMSO 20% (Sigma-Aldrich). For cell culture medium, RPMI medium was used to culture HeLa and MCF-7 cell line supplemented with Fetal Bovine Serum (FBS) and antibiotic-antimycotic.

2.2. Equipment

Equipment used in this study were moisture balance (Mettler Toledo™), heating and drying Oven (Memmert), Thin-Layer-Chromatography Chamber, ELISA Microplate reader, Bio-Safety Cabinet level 2, cell culture CO<sub>2</sub> incubator, and serological pipette.

19

2.3. Analytical Procedure

2.3.1. Sample Preparation and Extraction

Preparation of sample and extraction were performed based on previous study (Audah *et al.*, 2018). Sample preparation was carried out using an oven with a temperature of less than 50 °C until the samples were dry. The time needed for leaves was approximately 3 days, while the roots and stems were about 6 days to dry. The dried samples were stored in plastic and the moisture was kept and observed.

Extraction method used was gradient maceration (Farida *et al.*, 2014). Four different solvents were used, which were *n*-hexane, ethyl acetate, ethanol and water. yields were then determined based on the ratio of concentrated extract weight with initial sample weight. Extractions were performed triplicate.

2.3.2. Water Content Determination

The procedure for determining the water content of dried samples was carried out using a moisture balance. Five grams stored in an oven at 105°C for 5 hours, then weighed. The difference of mass before and after heating is the value of the water content of sample (Sulamsi *et al.*, 2016).

2.3.3. Qualitative Phytochemical Analysis

Phytochemical tests in this study include qualitative tests of alkaloid, phenolic, triterpenoid/steroid, and hydroquinone (Harborne, 1987).

2.3.4. Brine Shrimp Lethality Test (BSLT)

A tip of eggshell's *Artemia salina* (50–100 mg) was put into a container that had been filled previously with seawater and had been aerated using an aerator. The eggs were hatched at a temperature range of 25–30° C with light exposure until it was ready for use which was 48 hours old. Solution of extract for the test was made with various concentrations: 1000, 500, 100, and 50 ppm then dissolved in sea water with a little addition of Tween-80. Larvae were inserted into the vial (10 larvae/vial) and incubated for 24 hours. The number of dead larvae was quantified in 3 repetitions, then the obtained data were processed using Probit analysis on SPSS version 16 software to obtain LC<sub>50</sub> value.

2.3.5. MTT Cytotoxicity Assay

MCF-7 and Hela cells line were cultured on 96-well plates in the media of Rosewell Park Memorial Institute (RPMI) with 100 µL medium per well containing approximately 5,000 cells per well. A total of 17 µL of active extract with a concentration of 500 ppm was added to the cultured cells, incubated for 48 hours in a 5% CO<sub>2</sub> incubator at 37° C. Next, 100 µL of tetrazolium monoculture salt (MTT) were added and incubated again for 4 hours in 5% CO<sub>2</sub> incubator. Cells that are still alive will react with MTT to form formazan which can be seen violet. The formed Formazan was dissolved in 96% ethanol. The absorbance reading was in a microplate reader, 16 LISA spectrophotometer at λ = 595 nm. The tests were carried out using 1× RPMI media as a blank, MCF-7 breast cancer cells and HeLa cervical cancer cells as a negative control, and doxorubicin as a positive control.

3. Results

The yield of extracts of the 16 samples were between 0.25% (highlighted green) to 21.18% (highlighted yellow) that belonged to ethyl acetate extract of root of *Rhizophora apiculata* and water extract of leaf of *Xylocarpus granatum* respectively as shown in Table 2. The data clearly showed that root and leaf extract using non-polar solvent (*n*-hexane) resulted in lower yield percentage compared to more polar solvents (root samples 74 and 85, leaf samples 79 and 87) (Zahra *et al.*, 2017). This indicated that root sample contains less nonpolar constituents compared to other parts of plants.

Table 2: Yield of extracts (%)\*

Sample Code	Yield (%)	Sample Code	Yields (%)	Sample Code	Yield (%)	Sample Code	Yields (%)
74Ea	0.25	79Et	2.48	85H	0.5	86Ea	1.79
75A	20.32	79H	1.04	85Ea	0.89	87A	19.58
75Et	7.09	80A	3.72	86A	21.18	87H	2.41
79Ea	2.36	84Et	7.42	86Et	6.94	88A	7.06

\*Solvents abbreviation: H = *n*-hexane, Ea = ethyl acetate, Et = ethanol, A = water.

Water content of sample (Table 3) were kept below 10% which is a suitable percentage for simplisia analysis according to the Indonesian Herbal Pharmacopeia (Department of Health of the Republic of Indonesia., 2009). The largest shrinkage of drying is found in the leaves of all parts of the plant, while the smallest drying shrinkage is found in the stem.

**Table 3:** water content determination

Samples	Part	Shrinkage (%)	Water Content (%)	Samples	Part	Shrinkage (%)	Water Content (%)
<i>R. apiculatae</i>	Root	62	7.36	<i>A. marina</i>	Leaf	77	9.43
<i>B. gymnorhiza</i>	Leaf	63	5.34		Root	71	-
<i>S. caseolaris</i>	Leaf	84	7.23	<i>X. granatum</i>	Leaf	80	8.98
	Leaf	71	4.78				
<i>R. mucronata</i>	Stem	47	4.85	<i>C. tagal</i>	Leaf	82	9.45

Each simplisia does not always produce a positive reaction to all tests. This shows that mangrove plants are quite rich in secondary metabolites and the composition is not the same one to another (Joel and Bhimba, 2013). Furthermore, the results of this test can be a simple assumption for continued anticancer screening. The following results of the dried sample phytochemical test were presented in Table 4.

**Table 4:** Phytochemical qualitative analysis of samples

Sample Code	Alkaloid			Phenolic					
	M	W	D	F	Sa	Ta	T	S	Q
74	-	-	-	+	+++	+++	-	-	-
75	-	-	-	+++	+++	+++	-	++	-
79	-	-	-	+	+++	+++	-	++	-
80	-	-	-	-	+++	+++	-	-	+
84	-	-	-	++	++	++	-	+	-
85	-	-	-	-	+	-	-	+	-
86	-	-	-	-	+++	+	-	+	-
87	+	+	+	+	+	++	-	+	-
88	-	-	-	++	-	+++	-	+	-

Description: M=Mayer, W=Wagner, D=Dragendorf, T=Triterpenoid, S=Steroid, Q=Quinon, F=Flavonoid, Sa= Saponin, Ta= Tanin

26

**Brine Shrimp lethality Test (BSLT)** is one of the initial methods to estimate the level of toxicity of a compound based on the lethal concentration value of 50% (LC<sub>50</sub>). The shrimp larvae have thin skin and a sensitive body. If there are foreign compounds in the environment, the compound will be easily absorbed into the body of the larva diffusion. The toxic environment due to foreign compound will terminate the shrimp larvae. Chemical compounds are potentially bioactive if they have an LC<sub>50</sub> of less than 1000 ppm. In this study, there were 43 extracts (table 5) which have the potential to have toxic compounds. After obtaining the LC<sub>50</sub> value, some extracts were then selected which had LC<sub>50</sub> below 500 ppm (green highlighted) and some extracts which had LC<sub>50</sub> values more than 1000 ppm (yellow highlighted) for cytotoxicity testing. The lowest LC<sub>50</sub> was 40.45 ppm (87 H).

**Table 5:** Brine Shrimp Lethality Test LC<sub>50</sub>

Sample Code	LC <sub>50</sub> (ppm)	Sample Code	LC <sub>50</sub> (ppm)	Sample Code	LC <sub>50</sub> (ppm)	Sample Code	LC <sub>50</sub> (ppm)
74Ea	200.78	79Et	498.28	85H	499.23	86Ea	64.02

75A	>1000	79H	160.43	85Ea	>1000	87A	>1000
75Et	291.61	80A	>1000	86A	408.24	87H	40.45
79Ea	224.45	84Et	229.77	86Et	74.87	88A	488.93

The cytotoxicity of 16 extracts selected from the results of toxicity tests showed water extract of *Sonneratia caseolaris* leaf (88A), ethanol acetate extract of *Xylocarpus granatum* leaf (86Ea), and ethanol acetate extract of *Avicennia marina* root (85Ea) had a high percentage of inhibition in both cells. However, ethanol acetate extract of *Avicennia marina* root (85Ea) did not show bioactive potential that was toxic in BSLT because the obtained LC<sub>50</sub> obtained was below 1000 ppm. Meanwhile, ethanol extract of *Bruguiera gymnorrhiza* leaf (75Et) and ethanol acetate extract of *Rhizophora apiculata* root (74Ea) have a high percentage of inhibition in HeLa cells line, although it was low in MCF-7 cells line.

**Table 6: MTT Cytotoxicity Assay**

Sample Code	% Inhibition		LC <sub>50</sub> (ppm)	Sample Code	% Inhibition		LC <sub>50</sub> (ppm)
	MCF-7 Cells	HeLa Cells			MCF-7 Cells	HeLa Cells	
<b>74Ea</b>	<b>29.57</b>	<b>85.92</b>	<b>200.78</b>	<b>85Ea</b>	<b>98.06</b>	<b>94.04</b>	<b>&gt;1000</b>
75A	45.02	52.8	>1000	86A	44.4	64.8	408.24
<b>75Et</b>	<b>16.2</b>	<b>90.79</b>	<b>291.61</b>	86Et	74.87	85.92	351.52
79Ea	2.03	47.92	224.45	<b>86Ea</b>	<b>96.65</b>	<b>92.96</b>	<b>64.02</b>
79Et	18.09	51.71	498.28	87A	41.95	68.05	>1000
79H	-22.96	24.46	160.43	87H	52.53	29.06	40.45
80A	38.78	27.98	>1000	<b>88A</b>	<b>97.4</b>	<b>92.42</b>	<b>488.93</b>
84Et	48.37	68.5	229.77	Doxorubicin	94	95.76	-
85H	32.12	29.87	499.23	Control Cell	-	-	-

Description: **Green**= Highly potential, **Yellow**= slightly potential

The results of phytochemical analysis of *Avicennia marina* root (85Ea) indicated that there were only two active compounds, namely Saponin and Steroid which were detected in low-concentration. This result can explain the high value of LC<sub>50</sub> (>1000 ppm) of the extract which was not toxic enough for *Artemia salina*. Extracts of plants or animal are classified a cytotoxic when the LC<sub>50</sub> value <1000 ppm. An extract is considered very toxic when it has a LC<sub>50</sub> value < 30 ppm, and toxic if it has a LC<sub>50</sub> value 30-1000 ppm and is considered non-toxic if LC<sub>50</sub> > 1000 ppm. While pure compound was declared toxic if it has a value of LC<sub>50</sub> <200 ppm (Meyer *et al.*, 1982). Meanwhile, other extracts that have more than ten compounds were toxic for *Artemia salina*. For example, the extract of *Ceriops Tagal* leaf (87H) showed that there were seven compounds detected in the extract and had the lowest LC<sub>50</sub> value of all (40.45 ppm).

Moreover, the high inhibition percentage of ethanol acetate extract of *Avicennia marina* root (85Ea) in both cells line indicated that there was a possibility for false positive result. This is due the damaged mitochondria may be still able to reduce MTT to formazan crystals (Mosmann, 1983). Loveland *et al.* (1992) showed that cells with inactivated mitochondria were also able to produce formazan crystals as well as cells with active mitochondria. In addition, some ingredients of mangrove may have reducing activity such as the dehydrogenase activity of the cells that converts the MTT compound into the colored formazan. Therefore, the MTT assay may not be a suitable assay for mangrove extracts and it is important to conduct other cytotoxicity assay such as ATP assay, which gives more sensitive and reliable assay due to its interference-free feature (Karakas *et al.*, 2017).

Extracts of mangrove plant are considered to have potent cytotoxic activity against HeLa and MCF-7 cell lines. When compared between the cell lines HeLa cell lines are more suppressed than MCF-7 cell lines in general (Pradesh 18). The presence of different bio active compounds is the reason for its anticancer activity. Such compounds with antioxidant activity can inhibit mutation and cancer because

they scavenge free radicals or induce antioxidant enzymes (Karami *et al.*, 2012). For example, saponins that contained in almost all extracts can inhibit the replication of cellular DNA and the proliferation of cancer cells can be prevented (Yildirim and Kutlu, 2015). In addition, Saponins may stimulate the disintegration of the microtubular network or actin filaments of cancer cells, which can lead to further non-apoptotic cell death (Chen *et al.*, 2009).

#### 4. Conclusion

The methods and the results presented in this study suggested that developing extract library is a highly feasible and crucial for development of anticancer drugs discovery. Mangrove extracts are very potential as natural anticancer and antiproliferative agents. This report can be used as stepping stone towards acceleration anticancer drugs discovery.

#### 10 knowledgements

Authors would like to thank the Ministry of Research, Technology and Higher Education, Republic of Indonesia for the research grant to Kholis Abdurachim Au under contract Number: AGMT/Rec/A0018/III/2018. Authors also thank the Swiss German University for financial support through the Central Research Fund and the Faculty Research Fund.

#### References

- Audah, K. A., Septiyorini, D, Darmawan, A. G., Himawan, S and Batubara, I. (2018). Methods Development for Mangrove Extraction towards Standardization of Indonesian Extract Library Kholis. *Hayati Journal of Biosciences*. (unpublished manuscript).
- Audah, K.A., J Amsyir, F Almasyhur, A M Hapsari and H Sutanto (2018). Development of extract library from Indonesian biodiversity: exploration of antibacterial activity of mangrove *Bruguiera cylindrica* leaf extracts. IOP Conf. Ser.: Earth Environ. Sci. 130 012025.
- Bandaranayake, W.M. Bioactive compounds and Chemicals Constituents of Mangrove Plants, *Wetland Ecology and Management*, 10 (2002), pp. 421-452.
- Bray, F., Ferlay, J., & Soerjomataram, I. (2018). Global Cancer Statistics 2018 : GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries, *00(00)*, 1–31.
- Department of Health of the Republic of Indonesia. Indonesian Herbal Pharmacopeia. First Edition. (2009).
- Farida H, Severin JA, Gasem MH, et al. Nasopharyngeal carriage of *Streptococcus pneumoniae* in pneumonia-prone age groups in Semarang, Java Island, Indonesia. *PLOS One* 9(1) (2014), pp. 16–18.
- Giri, C., E. Ochieng, L. Tieszen, Z. Zhu, A. Singh, T. Loveland, J. Masek, and N. Duke (2011), Status and distribution of mangrove forests of the world using earth observation satellite data, *Global Ecology and Biogeography*, 20(1), 154-159.
- Harborne, J. B. (1987) 'Koleksi Buku 1987 Harborne, J. B. " Metode fitokimia : penuntun cara modern menganalisis tumbuhan / J. B. Harborne ; penerjemah Kosasih Padmawinata Iwang'
- H.-F. Chen, G.-H. Wang, Q. Luo, N.-L. Wang, and X.-S. Yao, "Two new steroidal saponins from *Allium macrostemon bunge* and their cytotoxicity on different cancer cell lines," *Molecules*, vol. 14, no. 6.
- Isil Yildirim and Turkan Kutlu, 2015. Anticancer Agents: Saponin and Tannin. *International Journal of Biological Chemistry*, 9: 332-340.
- Joel, E. L. and Bhimba, B. V. 'Evaluation of secondary metabolites from mangrove associated fungi *Meyerozyma guilliermondii*', *Alexandria Journal of Medicine*. 49(3) (2013), pp. 189–194.
- Karakas, D., Ari, F., & Ulukaya, E. (2017). The MTT viability assay yields strikingly false-positive viabilities although the cells are killed by some plant extracts, 919–925. <https://doi.org/10.3906/biy-1703-104>
- Karami, L., Majd, A., Mehrabian, S., Nabiuni, M., & Salehi, M. (2012). Antimutagenic and anticancer effects of *Avicennia marina* leaf extract on *Salmonella typhimurium* TA100 bacterium and human promyelocytic leukaemia HL-60 cells, 38, 349–355. <https://doi.org/10.2306/scienceasia1513-1874.2012.38.349>
- Kerry R.G., Pradhan P., Das G., Gouda S., Swamy M.K., Patra J.K. (2018) Anticancer Potential of Mangrove Plants: Neglected Plant Species of the Marine Ecosystem. In: Akhtar M., Swamy M. (eds) *Anticancer plants: Properties and Application*. Springer, Singapore
- Loveland BE, Johns TG, Mackay IR, Vaillant F, Wang ZX, HertzogPJ (1992). Validation of the MTT dye assay for enumeration of cells in proliferative and antiproliferative assays. *Biochem Int* 27: 501-510.
- Meyer BN, Ferrign RN, Putnam JE, Jacobson JE, Nicholas DE, McLaughlin JL, Brineshrimp: a convenient general bioassay for active plant constituents. *Planta Medica*, 1982, 45, 31-34
- Molnár, J. (2015). *Characterization of antiproliferative activities of triazolyl-steroid derivatives*.

- Mosmann T (1983). Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* 65: 55-63.
- Rintelen, K. Von, Arida, E., & Häuser, C. (2017). A review of biodiversity-related issues and challenges in megadiverse Indonesia and other Southeast Asian countries.
- Pradesh, A. (2018). Evaluation of In vitro Anticancer activity of aerial parts of *Avicennia alba* plant metholic extract against HeLa and MCF7 cell lines, 248–252.
- Sulasm ES, Indriwati SE, Suarsini E, 2016. Preparation of Various Type of Medicinal Plants Simplicia as Material of Jamu Herbal. International conference on education 2016, p1014-1024.
- Zahra, S. S., Ahmed, M., Qasim, M., Gul, B., Zia, M., & Mirza, B. (2017). Polarity based characterization of biologically active extracts of *Ajuga bracteosa* Wall . ex Benth . and RP-HPLC analysis, 1–16



# Iconiet-2018

---

## ORIGINALITY REPORT

---

<b>17</b> %	<b>17</b> %	<b>15</b> %	<b>5</b> %
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

---

## PRIMARY SOURCES

---

<b>1</b>	<b>journals.tubitak.gov.tr</b> Internet Source	<b>3</b> %
<b>2</b>	<b>ecos.watts.net.nz</b> Internet Source	<b>2</b> %
<b>3</b>	<b>www.japsonline.com</b> Internet Source	<b>2</b> %
<b>4</b>	<b>jeccr.biomedcentral.com</b> Internet Source	<b>1</b> %
<b>5</b>	<b>www.ncbi.nlm.nih.gov</b> Internet Source	<b>1</b> %
<b>6</b>	<b>worldwidescience.org</b> Internet Source	<b>1</b> %
<b>7</b>	<b>www.tandfonline.com</b> Internet Source	<b>1</b> %
<b>8</b>	<b>legacy.biotrop.org</b> Internet Source	<b>1</b> %
<b>9</b>	<b>Submitted to Institut Pertanian Bogor</b> Student Paper	<b>1</b> %

---

10	<a href="http://aip.scitation.org">aip.scitation.org</a> Internet Source	1%
11	<a href="http://www.thaiscience.info">www.thaiscience.info</a> Internet Source	1%
12	<a href="http://www.researchsquare.com">www.researchsquare.com</a> Internet Source	1%
13	L Sulmartiwi, D Y Pujiastuti, W Tjahjaningsih, Jariyah. " Potential of mangrove extract as an antioxidant agent using multilevel extraction ", IOP Conference Series: Earth and Environmental Science, 2018 Publication	<1%
14	<a href="http://onlinelibrary.wiley.com">onlinelibrary.wiley.com</a> Internet Source	<1%
15	<a href="http://idoc.pub">idoc.pub</a> Internet Source	<1%
16	<a href="http://pubs.rsc.org">pubs.rsc.org</a> Internet Source	<1%
17	<a href="http://patentscope.wipo.int">patentscope.wipo.int</a> Internet Source	<1%
18	<a href="http://scialert.net">scialert.net</a> Internet Source	<1%
19	<a href="http://link.springer.com">link.springer.com</a> Internet Source	<1%

20	<a href="http://ejournal.unsrat.ac.id">ejournal.unsrat.ac.id</a> Internet Source	<1%
21	<a href="http://journal.uinjkt.ac.id">journal.uinjkt.ac.id</a> Internet Source	<1%
22	<a href="http://repository.nwu.ac.za">repository.nwu.ac.za</a> Internet Source	<1%
23	Nur Imaniati Sumantri, Kenny Lischer, Dian Rachma Wijayanti, Tomy Abuzairi. "In silico study on RNA structures of intronic mutations of beta-globin gene", F1000Research, 2020 Publication	<1%
24	<a href="http://www.mdpi.com">www.mdpi.com</a> Internet Source	<1%
25	<a href="http://pdfs.semanticscholar.org">pdfs.semanticscholar.org</a> Internet Source	<1%
26	Indriati Ramadhani, Diva Larissa, Yeni Yuliani, Mellova Amir, Kusmiati Kusmiati. "Extraction, characterization, and biological toxicity of $\beta$ -glucans from <i>Saccharomyces cerevisiae</i> isolated from ragi", Journal of Microbial Systematics and Biotechnology, 2020 Publication	<1%

Exclude bibliography  On