

Screening of Antibacterial Compounds Against Escherichia coli from Hanjeli Seeds (Coix lacryma-jobi) Based on Metabolomics

by Kholis Audah

Submission date: 02-Jun-2023 10:56AM (UTC+0700)

Submission ID: 2107185801

File name: 4-TJNPR-2023-M090_Galley_Proof-CI.....pdf (304.15K)

Word count: 5975

Character count: 31687

Screening of Antibacterial Compounds Against *Escherichia coli* from Hanjeli Seeds (*Coix lacryma-jobi*) Based on MetabolomicsLa Ode Sumarlin^{1*}, Rudi Heryanto², Firman Riansyah³, Meyliana Wulandari¹, Kholis A. Audah⁴, Karimatul S. Irsyad²¹Department of Chemistry, Faculty of Science and Technology, UIN Syarif Hidayatullah Jakarta, Banten, 15412, Indonesia²Department of Chemistry, Faculty of Mathematics and Natural Sciences, IPB University, Bogor, 16680, Indonesia³Department of Mathematics Education, Islamic State Institute of Kendari, 93116, Indonesia⁴Department of Biomedical Engineering, Swiss German University, Tangerang, 15143, Indonesia

ARTICLE INFO

ABSTRACT

Article history:

Received 21 February 2023

Revised 15 March 2023

Accepted 02 May 2023

Published online 01 June 2023

Copyright: © 2023 Sumarlin *et al.* This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Hanjeli seed (*Coix lacryma-jobi* L.) is one of the important plants used as raw materials for food and traditional medicine. In addition, hanjeli seed also has several biological activities, one of which is antibacterial activity comes from the activity of the metabolites contained in the extract of the seed. This study aims to determine the antibacterial activity of hanjeli seed extract and identify the metabolites contained in some selected extracts using a metabolomics approach. Hanjeli seed was extracted with distilled water, 50% ethanol, 96% ethanol, and n-hexane. The crude extracts of hanjeli seed were evaluated for the activity against *Escherichia coli* ATCC 25922 and the metabolites were identified using liquid chromatography-mass spectrometry/mass spectrometry (LC-MS/MS). Hanjeli seed extracts had antibacterial activity ranging from 7.85±4.48 to 27.99±6.02%, with 50% ethanol extract had the highest antibacterial activity and n-hexane extract had the lowest. LC-MS/MS analysis of the 50% ethanol extract obtained 22 metabolites which were putatively identified. The antibacterial activity of the hanjeli seed extract is predicted to be caused by the phenolic compounds, such as gluconic acid, quinic acid, malic acid, citric acid, p-coumaric acid, and ferulic acid. The phenolic compounds contained in the hanjeli seed extract can inhibit the growth of *E. coli*, so that the extract can be applied as an antibacterial against *E. coli*.

Keywords: antibacterial, *Coix lacryma-jobi*, *E. coli*, LC-MS/MS, PCA

Introduction

The use of plants as traditional medicine has been used for a long time to treat health problems. Apart from being caused by the tendency of people in the current era to adopt a back-to-nature lifestyle, people realize that plants as medicines have advantages. Namely, they have few side effects compared to chemical treatments, are easy to find, economical, and can be simply made.¹

Herbal medicines have been widely used worldwide, both in developed and developing countries. Specifically for developing countries, the use of herbal medicines will reduce dependence on imports which are expensive and have side effects.² About 63% of the world's population uses traditional medicine.³ Regarding biodiversity, Indonesia ranks second after the Amazon forest, having many native plants with high potential for medicinal and pharmaceutical opportunities.⁴ Therefore, according to Batubara and Prastya (2020), further research on Indonesian medicinal plants must be carried out to provide holistic knowledge about the development of cosmetic and health products, for example, the human's oral health.⁵

One of the medicinal plants that can be used as traditional medicine is hanjeli (*Coix lacryma-jobi*). Besides being used as food, animal feed, and handicrafts, people in Asian countries also use hanjeli for traditional medicine.^{6,7}

This plant has been traditionally applied for treating diuretic, anti-rheumatic, antispasmodic, anti-inflammatory, anti-diarrheal, anthelmintic, antipyretic, antispasmodic, diuretic, hypoglycaemic, antioxidant, antifungal, and anti-cancer.⁸⁻¹² The part of the hanjeli plant that has antibacterial properties is the seeds. Compounds with antibacterial activity can inhibit bacterial growth by interfering with their metabolism.¹³ Zhu (2017) and Devaraj *et al.* (2020) reported that the chemical components identified in hanjeli seeds were alkaloids, tannins, saponins, flavonoids, triterpenoids, and phenols.^{14,15} According to Seukep *et al.* (2023), these compounds can function as antibacterials.¹⁶

Hanjeli is one of the antibacterial plants that has not been studied scientifically, so the information on the content that has antibacterial pharmacological activity in hanjeli seeds has not been known. Given that hanjeli contains antibacterial compounds and can be used as medicine, it is necessary to do the measurement of the antibacterial activity of hanjeli extract. In this study, we evaluated the activity of hanjeli seed extract against *E. coli* bacteria because *E. coli* is a pathogenic bacterium that often causes food spoilage and when consumed can cause mouth ulcers and gastroenteritis.¹⁷ Therefore, information about the activity of hanjeli seed extract against *E. coli* can be used as a reference in medical science, food safety, and environmental protection.

In general, the bioactive components of natural products are identified using the bioassay-guided isolation method, which is an identification method by repeatedly isolating compounds based on bioactivity until pure compounds are obtained. That method is carried out repeatedly so that it is less effective in terms of time and cost. An alternative method is needed to identify bioactive components using a metabolomics approach. The metabolomics approach is carried out by comprehensively extracting samples using a combination of solvents at different polarity levels.¹⁸ Then, the chemical content in the sample was identified using liquid chromatography-mass spectrometry/mass spectrometry (LC-MS/MS).

*Corresponding author. E mail: sumarlin@uinjkt.ac.id
Tel: +62-21) 7493606

Citation: Sumarlin LO, Heryanto R, Riansyah F, Wulandari M, Audah KA, Irsyad KS. Screening of Antibacterial Compounds Against *Escherichia coli* from Hanjeli Seeds (*Coix lacryma-jobi*) Based on Metabolomics. Trop J Nat Prod Res. 2023; 7(5):2867-2872 <http://www.doi.org/10.26538/tjnpr/v7i5.4>

Official Journal of Natural Product Research Group, Faculty of Pharmacy, University of Benin, Benin City, Nigeria.

The data obtained from the LC-MS/MS analysis is a complex data set, so the interpretation requires the chemometric methods. Chemometrics is a branch of science that derives data by applying mathematical and statistical methods to extract useful information from physical and chemical phenomena involved in a manufacturing process. Chemometrics is used for multivariate data collection and analysis protocols, calibration, process modeling, pattern recognition and classification, signal correction and compression, and statistical process control.⁴⁷ The chemometric pattern recognition technique used in this study is principal component analysis (PCA) because it can facilitate visualization in grouping data, evaluating similarities between groups or classes, and finding factors or reasons behind observed patterns through correlations based on the chemical or physicochemical properties of the samples.²⁰ In this study, the antibacterial compounds in hanjeli seed extract were predicted using LC-MS/MS-based metabolomic. LC-MS/MS results were further analyzed by PCA to determine the grouping pattern between the selected extracts with and without the addition of *E. coli* bacteria.

Materials and Methods

The materials used were hanjeli seeds (*Coix lacryma-jobi*), ethanol (Merck, Germany), n-hexane (Merck, Germany), DMSO (Merck, Germany), tetracycline (Darya-Varia, Indonesia), filter paper, methanol LC-MS grade (Merck, Darmstadt, Germany), acetone, distilled water LC-MS grade (Merck, Darmstadt, Germany), acetonitrile LC grade (Merck, Darmstadt, Germany), formic acid, distilled water, nutrient agar media, nutrient broth media, and *E. coli* ATCC 25922 bacteria obtained from the Laboratory of the Center for Tropical Biopharmaceutical Studies, LPPM IPB.

Extraction of Hanjeli Seeds.

Hanjeli seeds were obtained from the Medicinal Plants Cultivation and Conservation Unit, Tropical Biopharmaca Research Center, IPB University on 20 February 2022. Identification of the sample was performed by Mr. Taopik Ridwan (Trop-BRC, IPB University) with voucher specimen number BMK0336122016. The seeds were crushed into powder and extracted using the maceration method.²¹ A total of 50 g of hanjeli seed powder soaked in four different solvents at a ratio of 1:10 for 7x24. The solvents used were distilled water, 50% ethanol, 96% ethanol, and n-hexane. For solvent distilled water, maceration was carried out for 4 hours at 70 °C. The mixture was filtered and the filtrate was concentrated using a rotary evaporator.

Bacterial Inoculation

Inoculation of *E. coli* ATCC 25922 was done using NB (Nutrient Broth) and NA (Nutrient Agar) media. One dose (10 µL) of bacteria was added to 50 mL of NB medium and stirred thoroughly. The NB medium was incubated at 37°C for 24 hours. The turbidity of the NB medium indicated bacterial growth.

Antibacterial Activity Test with Disc Method

A total of 300 µL of bacterial suspension was added to 300 mL of NA medium. After mixing, the media was poured into 12 petri dishes. After becoming agar, five paper discs were added to each medium containing *E. coli* ATCC 25922. Test solutions with extract concentrations of 7%, positive control (tetracycline), and negative control (20% DMSO) was added to the disc paper. The plates were then incubated at 37 °C for 24 hours. The clear zone around the disc paper indicates the antibacterial activity of the hanjeli seed extract. The antibacterial activity of the extract is expressed in % inhibition and is calculated using the following equation:

$$\% \text{ inhibition} = 1 - \left(\frac{\text{positive control diameter} - \text{diameter of extract}}{\text{positive control diameter}} \right) \times 100\%$$

Sample Preparation

The extract with the highest antibacterial activity was analyzed using LC-MS/MS. The samples were divided into 3 groups, i.e. (I) hanjeli seed extract, (II) hanjeli seed extract in NB medium, and (III) hanjeli seed extract in NB media with the addition of *E. coli*. Group III was prepared by adding 1 mL of hanjeli seed extract to 2.8 mL of NB

medium and 0.2 mL of *E. coli* suspension. The mixture was incubated for 24 hours at 37 °C. The mixture was then centrifuged at 7000 rpm for 5 minutes. Acetonitrile (threefold of the supernatant volume) was added to the mixture and centrifuged at 12000 rpm for 1 minute. The supernatant obtained was analyzed using LC-MS/MS. Group II was prepared using the same method, but without the addition of bacterial suspension. Each group was prepared in three replicates. The second and third groups were compared to predict which compounds had antibacterial activity.

Identification of compounds with LC-MS/MS

The chemical compounds of hanjeli seed extract which have the highest antibacterial activity were analyzed using LC-MS/MS referring to the previous method with modifications.²² The extract solution was filtered with 0.22 µm filter membrane, and 5.0 µL of the filtrate was injected in LC-MS/MS. Samples were analyzed using two mobile phases, 0.1% formic acid in water (A) and 0.1% formic acid in acetonitrile (B), with a flow rate of 0.2 mL/min. The elution system used as a gradient, with the following composition of the mobile phase: 0-0.8 minutes (5%B), 0.8-12.4 minutes (5-95%B), 12-13 minutes (95%B), 13-15 minutes (5%B). The column used was Accucore™ Phenyl Hexyl (100 × 2.1 mm, 2.6 µm). The MS ionization source is ESI(-) with a Q-Orbitrap mass analyzer. The m/z range is from 80-1200, with ionization energies of 18, 35 and 53 eV. Other MS system parameters are set as follows: capillary temperature 320 °C; spray voltage 3.8 kV; sheath and auxiliary 15 and 3 arbitrary units, respectively; and resolving power of 70,000 FWHM.

Data from LC-MS/MS analysis were processed using Compound Discoverer 3.2 (Thermo Scientific, Waltham, USA) with online databases from ChemSpider and mzCloud. The steps for identifying metabolites include the selected spectra stage, alignment retention time, detected unknown compounds, grouping unknown compounds, predicting composition, filling gaps, normalizing areas, and marking background compounds. Metabolites were detected with a maximum mass error limit of 15 ppm and a minimum intensity of 1000000. Metabolites that were not identified using the database were not included in Table 1.

Statistical analysis

Extraction yield and antibacterial activity determined in three replicates and expressed as mean ± standard deviation. Principal component analysis (PCA) was used to differentiate between 50% ethanol extract in NB media with and without the addition of *E. coli* bacteria. The PCA model was performed using MetaboAnalyst ver 5.0 (<https://www.metaboanalyst.ca/>) with a variable peak area of all detected metabolites.

Results and Discussion

Hanjeli seed extraction aims to separate the bioactive compounds contained in the seeds. The solvent used in this extraction is distilled water, 96% ethanol, 50% ethanol, and n-hexane. The use of polar, semi-polar, and non-polar solvents aims to extract all the metabolites contained in hanjeli seeds which have different polarities. Extraction was carried out by maceration method. This method was chosen because the process is easier, more efficient, and does not damage the thermolabile compounds. In addition, Agustien *et al.*²³ also reported that the content of metabolites such as flavonoids obtained from the maceration method (1.39%) was higher compared to the soxhlet method (0.81%) which used heat.²³

The extraction yield shows the number of compounds that extracted during the maceration process. The extraction yield of the four solvents is shown by Figure 1a. The 96% ethanol extract has the highest extraction yield, which is 2.59%. This shows that the compounds in hanjeli seeds more extracted in 96% ethanol solvents. Ethanol has two groups associated with polarity, namely polar hydroxyl groups and nonpolar alkyl groups, causing polar and nonpolar compounds to be extracted and obtain high yield.²⁴

Hanjeli seed extract has several biological activities, one of which is an antibacterial. The antibacterial activity of hanjeli seed extract was measured using *E. coli* ATCC 25922 with various solvents. Hanjeli seed

extract from the four solvents can inhibit *E. coli* bacteria with varying inhibition percentages (Figure 1b). Hanjeli extract with 50% ethanol solvent has more potential to inhibit *E. coli* activity compared to hanjeli extract using other solvents. This shows that the 50% ethanol extract contains bioactive compounds that have potential as antibacterial. In contrast, the 96% ethanol extract, which had the highest extraction yield produced a lower inhibition percentage. This can be predicted because the many metabolites contained in the 96% ethanol extract have an antagonistic effect between compounds which can reduce their antibacterial activity.

The metabolites contained in the 30% ethanol extract with and without the addition of *E. coli* were identified using an LC-MS/MS based metabolomics approach. The column used is Accucore™ Phenyl Hexyl, commonly used for aromatic analytes and relatively polar compounds. The column provides fast separation with high resolution without the high back pressure required by the particles and has good sensitivity. This column has a reversed phase so that nonpolar compounds will pass through the column for longer than more polar compounds. The polarity of the mobile phase will change systematically with time, resulting in compounds with different polarity levels being separated from each other so that a gradient elution system is chosen.

Basepeak chromatogram of 7% ethanol extract in NB medium with and without the addition of *E. coli* bacteria is shown in Figure 2. The chromatogram was processed and 369 metabolites were detected. The 369 metabolites were then identified using the ChemSpider and mzCloud databases. A total of 22 metabolites were putatively identified based on the database and confirmed using the MS2 spectrum (Table 1). Meanwhile, we did not include unknown metabolites in Table 1. Most of the putatively identified metabolites belong to the phenolic, flavonoid, and fatty acid groups. This is in accordance with the results reported by Huang *et al.*²⁵ that hanjeli seed extract contains phenolic and flavonoid compounds.




Figure 1: Extraction yield (a) and antibacterial activity (b) of hanjeli seeds

The 50% ethanol extract in NB media with and without the addition of *E. coli* had similar chromatograms and was difficult to distinguish. Therefore, to differentiate the two groups we used principal component analysis (PCA) with the peak areas of all detected metabolites as variables. PCA is a technique for reducing the dimensionality of a data set and improving interpretation by minimizing loss of information.²⁶ The PCA score plot showed that the samples could be divided into two groups, i.e. 50% ethanol extract in NB media with and without the addition of *E. coli* using PC1 and PC2 with a total PC of 83.7% (Figure 3). The PC value shows the variance that can be explained by the two main components is 83.7% of the total variance. The two groups shown on the score plot indicated differences in metabolites detected with and without the addition of *E. coli* bacteria.

Antibacterial compounds against *E. coli* from hanjeli seed extract predicted based on literature study. The potency of hanjeli seed extract as an antibacterial is caused by the bioactive metabolites contained in the extract. Inhibition of bacteria by bioactive metabolites in extracts generally occurs through the mechanism of cell wall damage in bacteria which causes the plasma membrane to rupture, cytoplasm to exit, substance transport to be disrupted, and metabolism to be inhibited so that bacterial growth is also inhibited and even die.




Figure 2: Basepeak chromatogram of hanjeli seed extract + NB media (A) and seed extract + NB media + *E. coli* bacteria (B) in negative ionization mode




Figure 3: PCA score plot of hanjeli seed extract + NB media (A) and seed extract + NB media + *E. coli* bacteria (B)

Table 1: Putative identified metabolites in 50% ethanol extract with media and *E. coli*

No.	Name	Formula	Error (ppm)	Calc. MW	RT (min)	MS2
1	Gluconic acid	C ₆ H ₁₂ O ₇	-4.88	196.0574	1.304	195, 177, 87, 75
2	D-(-)-Quinic acid	C ₇ H ₁₂ O ₆	-5.02	192.0624	1.32	191, 173, 131, 129
3	4-Hydroxy-L-threonine	C ₄ H ₉ NO ₄	1.26	135.0533	1.34	134, 116
4	Isocitric acid	C ₆ H ₈ O ₇	-5.07	192.026	1.405	191, 173, 129
5	D-(+)-Malic acid	C ₄ H ₆ O ₅	-9.1	134.0203	1.407	133, 115, 71
6	L-gamma-Glutamyl-L-leucine	C ₁₁ H ₂₀ N ₂ O ₅	-2.02	260.1367	1.489	259, 130, 114, 112
7	Citric acid	C ₆ H ₈ O ₇	-5.16	192.026	1.493	191, 147, 129
8	Methylmalonic acid	C ₄ H ₆ O ₄	-10.15	118.0254	1.499	117, 73
9	N(2)-succinyl-L-glutamic acid	C ₉ H ₁₃ NO ₇	-3.03	247.0685	1.501	246, 128, 115
10	2-methylcitric acid	C ₇ H ₁₀ O ₇	-4.56	206.0417	1.504	205, 187, 161
11	Pyroglutamylisoleucine	C ₁₁ H ₁₈ N ₂ O ₄	-2.41	242.1261	5.894	241, 197, 130, 82
12	4-Hydroxybenzaldehyde	C ₇ H ₆ O ₂	-10.2	122.0355	6.213	121, 93, 67
13	p-coumaric acid	C ₉ H ₈ O ₃	-7	164.0462	6.515	163, 119, 93
14	2,3-Dihydro-1-benzofuran-2-carboxylic acid	C ₉ H ₈ O ₃	-6.71	164.0462	6.686	163, 119
15	Ferulic acid	C ₁₀ H ₁₀ O ₄	-5.43	194.0569	7.582	193, 147, 133
16	Corchorifatty acid F	C ₁₈ H ₃₂ O ₅	-2.32	328.2242	8.28	327, 229, 171
17	Cinniferyl ferulate	C ₂₀ H ₂₀ O ₆	-2.4	356.1251	8.342	355, 337, 219, 175
18	(15Z)-9,12,13-Trihydroxy-15-octadecenoic acid	C ₁₈ H ₃₄ O ₅	-2.39	330.2398	8.536	329, 229, 171
19	(10E,12E)-9-hydroperoxyoctadeca-10,12-dienoic acid	C ₁₈ H ₃₂ O ₄	-2.21	312.2294	9.859	311, 275, 235
20	9,10-dihydroxy-12Z-octadecenoic acid	C ₁₈ H ₃₄ O ₄	-2.06	314.2451	10.249	313, 201, 171, 127
21	13S-hydroxyoctadecadienoic acid	C ₁₈ H ₃₂ O ₃	-2.58	296.2344	11.865	295, 183
22	2,2'-Methylenebis(4-methyl-6-tert-butylphenol)	C ₂₃ H ₃₂ O ₂	-1.92	340.2396	13.663	339, 163

Based on literature study, the antibacterial activity of hanjeli seed extract is ²⁷ed by phenolic compounds, i.e. phenolic acids and flavonoids. This is in ⁴ accordance with the research of Dzoyem *et al.*²⁷ who reported that in antibacterial activity (against Gram-positive and Gram-negative ⁴⁶acteria), DNA, RNA, and protein synthesis were ⁷ inhibited by three flavonoids isolated from *Dorstenia* species. In addition, Shamsudin *et al.*²⁸ also reported that the hydroxylation of C5, C7, C3', and C4' in the flavonoid structure can increase bacterial inhibition.

Research from Wu *et al.*²⁹ reported the mechanism of action of five flavonoids against *E. coli*. These compounds were effective via rigidifying the membrane of liposomal. The molecular hydrophobicity (C log P) and charges on the C atom at position three ² might play a role in the intercalation of liposomal model membranes. He *et al.*³⁰ screened the antimicrobial mechanism of flavonoids (kaempferol, hesperetin) for inhibitory activity against *E. coli* through the cell membranes and liposomal model. The studies result that interaction between the group of polar heads of the membrane model and the hydrophobic regions may damage the *E. coli* membrane.

Furthermore, Liu *et al.*³¹ reported the mechanism of bacterial inhibition by ¹ phenolic acid compounds. Phenolic acid can inhibit bacterial growth by destabilizing the bacterial cytoplasmic membrane, changing the permeability of the bacterial plasma membrane, inhibiting extracellular microbial enzymes, directly changing microbial metabolism, changing the physicochemical surface properties of bacteria, and removing

microbes from the substrate needed for bacterial growth. In addition, phenolic acid can also change the polarity of ¹acteria by changing the surface electron acceptor of bacteria in gram-positive (increased acceptor component) and gram-negative (decreased acceptor component) strains.³²

Several metabolites identified in hanjeli seed extract ¹³ have been reported to have inhibitory activity against *E. coli* bacteria, such as gluconic acid, quinic acid, malic acid, citric acid, p-coumaric acid, and ferulic acid. Masoura *et al.*³³ reported that there was a synergistic effect between gluconic acid and H₂O₂ in inhibiting the growth of *E. coli* bacteria. This synergistic effect causes membrane depolarization, cell wall destruction, and finally inhibition of *E. coli* growth.

Quinic acid and citric acid are short-chain non-aromatic hydroxy acids which have important roles in plant metabolism. Both of these compounds have been reported ² have antibacterial activity. Quinic acid and citric acid can inhibit the growth of *E. coli* bacteria with an MIC of 500 µg/mL.³⁴ Quinic acid has the potential as an antibacterial by damaging ¹¹ membranes and interfering with cell metabolic activity.³⁵ Bai *et al.*³⁶ reported ²¹ quinic acid can inhibit the growth of bacteria by interfering with the oxidative phosphorylation pathway, and changing levels of glycerophospholipids and fatty acids to disrupt membrane ¹ fluidity. After penetrating the cell membrane, quinic acid affects ribosome function and aminoacyl-tRNA synthesis, thereby interfering with protein synthesis to inhibit bacterial growth.

Meanwhile, citric acid can inhibit bacterial growth by passing through the microbial membrane. Once inside the cytoplasm, it dissociates into citric acid anions and generates protons which cause acidification of the intracellular medium while causing functional and structural damage to the bacterial cell.³⁷ In addition, Kang *et al.*¹⁷ sorted that citric acid can pass through cell membranes and promote superoxide dismutase (SOD) inactivation in *E. coli* O157:H7. This induces reactive oxygen species (ROS) and promotes cell inactivation by activating lipid oxidation, causing cell membrane damage. Al-Rousan *et al.*³⁹ also reported that citric acid could inhibit the growth of *E. coli* bacteria in salads, thereby reducing the risk of contamination by *E. coli* bacteria.³⁹ Malic acid³⁷ included in organic acids that can act as an antibacterial caused by a decrease in pH below the growth range of microorganisms and inhibition of metabolism by con dissociation acid molecules. In addition, according to Massilia *et al.*⁴⁰ transmission electron microscopy showed that malic acid produced damage in the cell cytoplasm of pathogens without apparent changes in the cell membrane. Ballal *et al.*⁴¹ found that 17% of malic acid showed antimicrobial activity on *Enterococcus faecalis*, *Candida albicans*, and *Staphylococcus aureus*. The presence of malic acid in trimethoprim can ease its solubility and antibacterial activity.⁴² Ferulic acid and p-coumaric acid are phenolic acids which have been reported to have activity against *E. coli* bacteria. Borges *et al.*⁴³ reported MIC and MBC ferulic acid for *E. coli* was 100 µg/mL and 2500 µg/mL, respectively. Ferulic acid can cause damage to cell membranes, thereby inhibiting bacterial growth. These compounds damage cell membranes by changing the properties of the membrane through hydrophobic changes and a decrease in negative surface charge.⁴⁴ While p-coumaric acid can inhibit bacterial growth by disrupting cell membranes and binding to bacterial genomic DNA to inhibit cellular functions.⁴⁵

Conclusion

Hanjeli seed extract inhibit the activity of *E. coli* bacteria, with the highest percentage of inhibition found in the extract with 50% ethanol solvent. A total of 22 compounds were identified in hanjeli seed extract. 44 identified compounds belong to the phenolic and fatty acid groups. The identified phenolic compounds are predicted to influence the antimicrobial activity of hanjeli seed extract.

Conflict of Interest

The authors declare no conflict of interest.

Authors' Declaration

The authors hereby declare that the work presented in this article is original and that any liability for claims relating to the content of this article will be borne by them.

Acknowledgements

This study was funded by the Institute for Research and Community Service (LP2M) through the Research and Publishing Centre, UIN Syarif Hidayatullah Jakarta in the Fiscal Year 2022 through SK Un.01/KPA/223/2022. The authors are grateful to the Tropical Biopharmaca Research Center, IPB University, for providing all facilities to conduct this study.

References

- Rahayu YYS, Araki T, Rosleine D. Factors affecting the use of herbal medicines in the universal health coverage system in Indonesia. *J Ethnopharmacol.* 2020; 260: 112974.
- Sam S. Importance and effectiveness of herbal medicines. *J. Pharmacogn Phytochem.* 2019; 8(2): 354-357.
- [WHO] World Health Organization. 2023. WHO Traditional Medicine Strategy 2014-2023. Geneva (CH): WHO Pr.

- Yaman E, Woerdenbag HJ, Kayser O. Jamu: Indonesian traditional herbal medicine towards rational phytopharmacological use. *J Herb Med.* 2014; 4(2): 51-73
- Batubara I, Prasty ME. Potential use of Indonesian medicinal plants for cosmetic and oral health: A review. *Jurnal Kimia Valensi.* 2020; 6(1): 118-132.
- Feng L, Zhao Y, Zhang Z, Zhang S, Zhang H, Yu M, Ma Y. The edible and medicinal value of *Coix lacryma-jobi* and key cultivation techniques for high and stable yield. *Nat Resour.* 2020; 11: 569-575.
- Tsay GJ, Lin Y, Hsu C, Tang F, Kuo Y, Chao C. Adlay hull extracts attenuate β -amyloid-induced neurotoxicity and oxidative stress in PC12 cells through antioxidant, anti-inflammatory, and antiapoptotic activities. *Biochem Biophys Rep.* 2021; 26: 101020.
- Patel B, Patel G, Shah S, Parmar SA. Review: *Coix lacryma jobi* L. *Res J Pharmacognosy and Phytochem.* 2017; 9(4): 248-252.
- Ruan J, Weng W, Yan J, Zhou Y, Chen H, Ren M, Cheng J. *Coix lacryma-jobi* chymotrypsin inhibitor displays antifungal activity. *Pestic Biochem Physiol.* 2019; 160: 49-57.
- Yu Q, Ye G, Lei Z, Yang R, Chen R, He T, Huang S. An isolated compound from stems and leaves of *Coix lacryma-jobi* L. and its anticancer effect. *Food Biosci.* 2021; 42: 101047.
- Xu L, Gao S, Xu H, Wang X, Hou Y, Liang N, Chen X. Impact of incubation on nutritional and antioxidant properties of defatted adlay (*Coix lacryma-jobi* L.) bran. *LWT – Food Sci Technol.* 2021; 137: 110463.
- He W, Yin M, Yang R, Zhao W. Optimization of adlay (*Coix lacryma-jobi*) bran oil extraction: Variability in fatty acids profile and fatty acid synthase inhibitory activities. *Biocatal Agric Biotechnol.* 2020; 28: 101740.
- Shaik G, Sujatha N, Mehar SK. Medicinal plants as source of antibacterial agents to counter *Klebsiella pneumoniae*. *J Appl Pharm. Sci.* 2014; 4(1): 135-147.
- Zhu F. Coix: Chemical composition and health effects. *Trends Food Sci Technol.* 2017; 61: 160-175.
- Devaraj RD, Jeepipalli SPK, Xu B. Phytochemistry and health promoting effects of Job's tears (*Coix lacryma-jobi*) – A critical review. *Food Biosci.* 2020; 34: 100537.
- Seukep AJ, Mbuntcha HG, Zeuko'o EM, Woquan LS, Nembu NE, Bomba FT, Watching D, Kuete V. Established antibacterial drugs from plants. In: Kuete V. Advances in botanical research. Amsterdam: Elsevier; 2023. 81-149.
- Ma E, An Y, Zhang G, Zhao M, Iqbal MW, Zayed HM, Qi X. Enhancing the antibacterial activity of *Lactobacillus reuteri* against *Escherichia coli* by random mutagenesis and delineating its mechanism. *Food Biosci.* 2023; 51: 102209.
- Yuliana ND, Khatib A, Choi YH, Verpoorte R. Metabolomics for bioactivity assessment of natural products. *Phytother Res.* 2011; 25 (2): 157-169.
- Singh I, Juneja P, Kaur B, Kumar P. Pharmaceutical applications of chemometric techniques. *Int Sch Res Notices.* 2013; 795178.
- Shiyan S, Zubaidah, Pratiwi G. Chemometric approach to assess response correlation and its classification in simplex centroid design for pre-optimization stage of catechin-snedds. *Res J Pharm and Tech.* 2021; 14: 5863-5870.
- Susilawati E, Adnyana IK, Kusuma E. Aktivitas antidiabetes dari ekstrak etanol biji hanjeli (*Coix lacryma-jobi*) pada mencit galur swiss webster yang diinduksi aloksan. *Galenika J. Pharm.* 2017; 2(2): 77-82.
- Tang Y, Lou Z, Young L, Wang H. Screening of antimicrobial compounds against *Salmonella typhimurium* from burdock (*Arctium lappa*) leaf based on metabolomics. *Eur Food Res Technol.* 2015; 240: 1203-1209.
- Agustien GS, Susanti, Sucitra F. Effect of different extraction method on total flavonoid contents of *Sansevieria trifasciata* P. leaves extract. *Galenika J Pharm.* 2021; 7 (2): 143-150.

24. Sulaiman ISC, Basri M, Masoumi HRM, Chee WJ, Ashari SE, Ismail M. Effects of temperature, time, and solvent ratio on the extraction of phenolic compounds and the anti-radical activity of *Clinacanthus nutans* Lindau leaves by response surface methodology. *Chem Cent J*. 2017; 11: 54.
25. Huang DW, CH Wu, CK Shih, CY Liu, PH Shih, TM Shieh, SM Hsia. Application of the solvent extraction technique to investigation of the antiinflammatory activity of adlay bran. *Food Chem*. 2014; 145: 445–453.
26. Paul A, de Boves Harrington P. Chemometric applications in metabolomic studies using chromatography-mass spectrometry. *TrAC Trends Anal Chem*. 2021; 135: 116165.
27. Dzoyem JP, Hamamoto H, Ngameni B, Ngadjui BT, Sekimizu K. Antimicrobial action mechanism of flavonoids from *Dorstenia* species. *Drug Discov Ther*. 2013; 7(2): 66–72.
28. Shamsudin NF, Ahmed QU, Mahmood S, Ali Shah SA, Khatib A, Mukhtar S, Alsharif MA, Parveen H, Zakaria ZA. Antibacterial effects of flavonoids and their structure-activity relationship study: A comparative interpretation. *Molecules*. 2022; 27(4): 1149.
29. Wu T, He M, Zang X, Zhou Y, Qiu T, Pan S, Xu XA. Structure-activity relationship study of flavonoids inhibitors of *E. coli* by membrane interaction effect. *Biochim Biophys Acta Biomembr*. 2013; 1828(11): 2751–2756.
30. He M, Wu T, Pan S, Xu X. Antimicrobial mechanism of flavonoids against *Escherichia coli* ATCC 25922 by model membrane study. *Appl Surf Sci*. 2014; 305: 515–521.
31. Liu J, Du C, Beaman HT, Monroe MBB. Characterization of phenolic acid antimicrobial and antioxidant structure-property relationships. *Pharmaceutics*. 2020; 12(5): 419.
32. Borges A, Ferreira C, Saavedra MJ, Simões M. Antibacterial activity and mode of action of ferulic and gallic acids against pathogenic bacteria. *Microb Drug Resist*. 2013; 19: 256–265.
33. Masoura M, Passaretti P, Overton TW, Lund PA, Gkatzionis K. Use of a model to understand the synergies underlying the antibacterial mechanism of H₂O₂-producing honeys. *Sci Rep*. 2020; 10(1): 17692.
34. Adameczak A, Ożarowski M, Karpiński TM. Antibacterial activity of some flavonoids and organic acids widely distributed in plants. *J Clin Med*. 2019; 9(1): 109.
35. Bai J, Wu Y, Wang X, Liu X, Zhong K, Huang Y, Chen Y, Gao H. In vitro and in vivo characterization of the antibacterial activity and membrane damage mechanism of quinic acid against *Staphylococcus aureus*. *J Food Saf*. 2017; 38(1): e12416.
36. Bai J, Wu Y, Bu Q, Zhong K, Gao H. Comparative study on antibacterial mechanism of shikimic acid and quinic acid against *Staphylococcus aureus* through transcriptomic and metabolomic approaches. *Lwt-Food Sci Technol*. 2022; 153: 112441.
37. Burel C, Kala A, Purevdorj-Gage L. Impact of pH on citric acid antimicrobial activity against Gram-negative bacteria. *Lett Appl Microbiol*. 2021; 72(3): 332–340.
38. Kang JW, Lee HY, Kang DH. Synergistic bactericidal effect of hot water with citric acid against *Escherichia coli* O157:H7 biofilm formed on stainless steel. *Food Microbiol*. 2021; 95: 103676.
39. Al-Rousan WM, Olaimat AN, Osaili TM, Al-Nabulsi AA, Ajo RY, Holley RA. Use of acetic and citric acids to inhibit *Escherichia coli* O157:H7, *Salmonella Typhimurium* and *Staphylococcus aureus* in tabbouleh salad. *Food Microbiol*. 2018; 73: 61–66.
40. Massilia RMR, Melgar JM, Bellosso OM. Antimicrobial activity of malic acid against *Listeria monocytogenes*, *Salmonella Enteritidis* and *Escherichia coli* O157:H7 in apple, pear and melon juices. *Food Control*. 2009; 20(2): 105–112.
41. Ballal NV, Yegneswaran PP, Mala K, Bhat KS. In vitro antimicrobial activity of maleic acid and ethylenediaminetetraacetic acid on endodontic pathogens. *J Endod*. 2011; 112(5): 696–700.
42. Yuliandra Y, Hutabarat LJ, Ardila R, Octavia MD, Zaini E. Enhancing solubility and antibacterial activity using multi-component crystals of trimethoprim and malic acid. *Pharmacy Education*. 2021; 21(2): 296–304.
43. Borges A, Ferreira C, Saavedra MJ, Simões M. Antibacterial activity and mode of action of ferulic and gallic acids against pathogenic bacteria. *Microb Drug Resist*. 2013; 19(4): 256–65.
44. Almasaudi S. The antibacterial activities of honey. *Saudi J Biol Sci*. 2021; 28(4): 2188–2196.
45. Lou Z, Wang H, Rao S, Sun J, Ma C, Li J. p-Coumaric acid kills bacteria through dual damage mechanisms. *Food Control*. 2012; 25: 550–554.

Screening of Antibacterial Compounds Against Escherichia coli from Hanjeli Seeds (Coix lacryma-jobi) Based on Metabolomics

ORIGINALITY REPORT

20%

SIMILARITY INDEX

18%

INTERNET SOURCES

16%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	www.mdpi.com Internet Source	2%
2	eprints.mums.ac.ir Internet Source	2%
3	tjnpr.org Internet Source	1%
4	link.springer.com Internet Source	1%
5	www.hindawi.com Internet Source	1%
6	www.researchgate.net Internet Source	1%
7	mdpi-res.com Internet Source	1%
8	jurnal.ugm.ac.id Internet Source	1%

journal.uinjkt.ac.id

9	Internet Source	1 %
10	ti.fst.uinjkt.ac.id Internet Source	1 %
11	brief.land Internet Source	1 %
12	bio-protocol.org Internet Source	1 %
13	Mouming Zhao, Dashuai Zhu, Dongxiao Sun-Waterhouse, Guowan Su, Lianzhu Lin, Xiao Wang, Yi Dong. "In Vitro and In Vivo Studies on Adlay-Derived Seed Extracts: Phenolic Profiles, Antioxidant Activities, Serum Uric Acid Suppression, and Xanthine Oxidase Inhibitory Effects", Journal of Agricultural and Food Chemistry, 2014 Publication	1 %
14	Mari Davies, Prabha Peramuhendige, Lloyd King, Melanie Golding, Apoorva Kotian, Mark Penney, Syeda Shah, Nenad Manevski. "Evaluation of In Vitro Models for Assessment of Human Intestinal Metabolism in Drug Discovery", Drug Metabolism and Disposition, 2020 Publication	<1 %
15	rjptonline.org Internet Source	<1 %

16	biodiversitas.mipa.uns.ac.id Internet Source	<1 %
17	www.esp.org Internet Source	<1 %
18	journals.plos.org Internet Source	<1 %
19	moam.info Internet Source	<1 %
20	www.ncbi.nlm.nih.gov Internet Source	<1 %
21	Jinrong Bai, Yanping Wu, Qian Bu, Kai Zhong, Hong Gao. "Comparative study on antibacterial mechanism of shikimic acid and quinic acid against Staphylococcus aureus through transcriptomic and metabolomic approaches", LWT, 2022 Publication	<1 %
22	pdfs.semanticscholar.org Internet Source	<1 %
23	www.thefreelibrary.com Internet Source	<1 %
24	Anabela Borges, Carla Ferreira, Maria J. Saavedra, Manuel Simões. "Antibacterial Activity and Mode of Action of Ferulic and	<1 %

Gallic Acids Against Pathogenic Bacteria", Microbial Drug Resistance, 2013

Publication

- 25 Enyun Ma, Yingfeng An, Guoyan Zhang, Mei Zhao, Muhammad Waheed Iqbal, Hossain M. Zabed, Xianghui Qi. "Enhancing the antibacterial activity of *Lactobacillus reuteri* against *Escherichia coli* by random mutagenesis and delineating its mechanism", *Food Bioscience*, 2022

Publication

- 26 Mohamad Rafi, Dien Atin Boritnaban, Dewi Anggraini Septaningsih, Fifi Gus Dwiyaniti et al. "Untargeted metabolomics analysis of *Diospyros celebica* Bakh. from three different geographical origins in Sulawesi island using UHPLC-Q-Orbitrap HRMS", *Wood Science and Technology*, 2022

Publication

- 27 media.neliti.com

Internet Source

- 28 "Bioactive Factors and Processing Technology for Cereal Foods", Springer Science and Business Media LLC, 2019

Publication

- 29 chenglong yang, Xiuwen Ban, Mingqiang Zhou, Yu Zhou et al. "Construction of a High-Density Genetic Map Based on Large-Scale

Marker Development in *Coix lacryma-jobi* L . Using Specific-Locus Amplified Fragment Sequencing (SLAF-seq)", Cold Spring Harbor Laboratory, 2022

Publication

30

cmuj.cmu.ac.th

Internet Source

<1 %

31

doaj.org

Internet Source

<1 %

32

repositorio-aberto.up.pt

Internet Source

<1 %

33

rgd.mcw.edu

Internet Source

<1 %

34

www.huveta.hu

Internet Source

<1 %

35

Dongwei Liu, Mei Liu, Dehao Meng, Yihan Mu, Tao Wang, Zhaolin Lv. "Harsh Sensitivity and Mechanism Exploration of an Antibacterial Peptide Extracted from Walnut Oil Residue Derived from Agro-Industrial Waste", Journal of Agricultural and Food Chemistry, 2022

Publication

<1 %

36

Theophilus Fadjare Frempong, Mercy Badu, Vivian Etsiapa Boamah, Isaac Kingsley Amponsah et al. "Phenolic Content, Antioxidant Properties and Antimicrobial

<1 %

Activities of the Extracts from Funtumia africana and Funtumia elastica", Chemistry Africa, 2021

Publication

37 jurnal.permi.or.id <1 %
Internet Source

38 soft-tox.org <1 %
Internet Source

39 www.frontiersin.org <1 %
Internet Source

40 www.rasayanjournal.co.in <1 %
Internet Source

41 Gina Porras, François Chassagne, James T. Lyles, Lewis Marquez et al. "Ethnobotany and the Role of Plant Natural Products in Antibiotic Drug Discovery", Chemical Reviews, 2020
Publication

42 www.acarindex.com <1 %
Internet Source

43 "Plant and Human Health, Volume 2", Springer Science and Business Media LLC, 2019
Publication

44 Atiqur Rahman, Shah Alam Siddiqui, Feyza Oke-Altuntas, Sezer OKAY, Fatih GÜL, Ibrahim <1 %

Demirtas. "Phenolic Profile, Essential Oil Composition and Bioactivity of *Lasia spinosa* (L.) Thwaites", Brazilian Archives of Biology and Technology, 2019

Publication

45

D. Zhang, S.-F. Yu, Y. Ma, K. Xu et al. "Chemical Structure and Concentration of Intratumor Catabolites Determine Efficacy of Antibody Drug Conjugates", Drug Metabolism and Disposition, 2016

Publication

46

Faegheh Farhadi, Bahman Khameneh, Mehrdad Iranshahi, Milad Iranshahy. "Antibacterial activity of flavonoids and their structure-activity relationship: An update review", Phytotherapy Research, 2018

Publication

47

Inderbir Singh, Prateek Juneja, Birender Kaur, Pradeep Kumar. "Pharmaceutical Applications of Chemometric Techniques", ISRN Analytical Chemistry, 2013

Publication

48

Jane D. Fudyma, Jamee Lyon, Roya AminiTabrizi, Hans Gieschen et al. "Untargeted metabolomic profiling of reveals novel antimicrobial metabolites ", Plant Direct, 2019

Publication

<1 %

<1 %

<1 %

<1 %

49

Tahara, Kayoko, Yuka Yano, Kiyotada Kanagawa, Yoshihiro Abe, Junko Yamada, Shiro Iijima, Masataka Mochizuki, and Takashi Nishikawa. "Successful Preparation of Metabolite of Troglitazone by In-Flow Electrochemical Reaction on Coulometric Electrode", CHEMICAL & PHARMACEUTICAL BULLETIN, 2007.

Publication

<1 %

50

repositorio.ipen.br

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On