

REFERENCES

Ajo, R. Y. (2017). Application of hydrocolloids as coating films to reduce oil absorption in fried potato-chip-based pellets. *Pak. J. Nutr.*, 16, pp. 805-812

Altunakar, B., Sahin, S., & Sumnu, G. (2004). Functionality of Batters Containing Different Starch Types for Deep-fat Frying of Chicken Nuggets. *European Food Research and Technology* 218(4), pp. 318-322.

Anderson, A. K., Guraya, H. S., James C, & Salvaggio L. (2002). Digestibility and Pasting Properties of Rice Starch Heat-moisture Treated at the Melting Temperature (tm). *Starch/Stärke* 54, pp. 401–409.

Asp, N. G. (1994). Nutritional Classification and Analysis of Food Carbohydrates. *The American Journal of Clinical Nutrition* 59(3), pp. 679S-681S.

Badan Penelitian dan Pengembangan Kesehatan. (2013). *Laporan Hasil Riset Kesehatan Dasar (Riskesdas) 2013*. Jakarta, Indonesia: Kementerian Kesehatan RI.

BPS. (2015). *Statistik Indonesia 2015*. Jakarta, Indonesia: Badan Pusat Statistik.

Brown, W. H., Poon, T. (2015). *Introduction to organic chemistry (3rd 3d.)* Wiley. ISBN 0-471-44451-0.

Cahill, L. E., Pan, A., Chiuve, S. E., Sun, Q., Willett, W. C., Hu, F. B., & Rimm, E. B. (2014). Fried-food consumption and risk of type 2 diabetes and coronary artery disease: a prospective study in 2 cohorts of US women and men. *American Journal of Clinical Nutrition* 100(2), pp. 667-675.

Dana, D. & Saguy, I. (2006). Review: Mechanism of oil uptake during deep-fat frying and the surfactant effect-theory and myth. *Advances in Colloid and Interface Science*, 128-130, pp. 267-272.

Dogan, S. F., Sahin, S., & Sumnu, G. (2005). *Effects of Batters Containing Different Protein Types on The Quality of Deep-Fat-Fried Chicken Nuggets. European Food Research and Technology* 220, pp. 502–508.

Dreher, M., Dreher, C., Berry, J., & Fleming, S. (1984). Starch digestibility of foods: A nutritional perspective. *C R C Critical Reviews in Food Science and Nutrition* 20(1), pp. 47-71.

Evans, I. D., Haisman, D. R. (1982). The Effect of Solutes on the Gelatinization Temperature Range of Potato Starch. *Starch -Stärke*. 34 (7), pp. 224–231.

Faridah, D. N. (2011). *Perubahan Karakterisasi Kristalin Pati Garut (Maranta arundinaceae L.) dalam Pengembangan Pati Resisten Tipe III*. Bogor: Institut Pertanian Bogor.

Fausto, F. D., Kacchi, A. I., and Mehta, D. (1997). Starch Products in Confectionery. *Beverage Food World* 24(4), pp. 4-16.

Fiszman, S., & Salvador, A. (2003). Recent Developments in Coating Batters. *Trends in Food Science & Technology* 14, pp. 399–407.

Gadiraju, T. V., Patel, Y., Gaziano, J. M., & Djoussé. (2015). Fried Food Consumption and Cardiovascular Health: A Review of Current Evidence. *Nutrients* 7(10), pp. 8424-8430.

Lehmann, U., Jacobasch, G., & Schmiedl, D. (2002). Characterization of Resistant Starch Type 3 from Banana (*Musa acuminata*). *Journal of Agricultural and Food Chemistry* 50(18), pp. 5236-5240.

Mellema, M. (2003). Mechanism and Reduction of Fat Uptake in Deep-fat Fried Foods. *Trends in Food Science & Technology* 14, pp. 364-373.

Moongngarm, A. (2013). Chemical Compositions and Resistant Starch Content in Starchy Foods. *American Journal of Agricultural and Biological Sciences* 8(2), pp. 107-113.

Moreira, R. G., Sun, X., & Chen, Y. (1997). Factors Affecting Oil Uptake in Tortilla Chips in Deep-fat Frying. *Journal of Food Engineering* 31, pp. 485-498.

Moreira, R. G. & Barrufet, M. A. (1998). A New Approach to describe Oil Absorption in Fried Foods: A Simulation Study. *Journal of Food Engineering* 35(1), pp. 1-22.

Muchtadi, D., Palupi, N., S., & Astawan, M. (1992). *Petunjuk Laboratorium Metode Kimia Biokimia dan Biologi dalam Evaluasi Nilai Gizi Pangan Olahan*. Departemen Pendidikan dan Kebudayaan, Direktorat Jendral Pendidikan Tinggi, Pusat Antar Universitas, Institut Pertanian Bogor.

Musita, N. (2009). *Kajian Kandungan dan Karakteristik Pati Resisten dari Berbagai Varietas Pisang*. Balai Riset dan Standarisasi Industry Bandar Lampung. *Jurnal Teknologi Industri dan Hasil Pertanian* 14(1), pp. 68-79.

Nugent, A. P. (2005). Health Properties of Resistant Starch. *British Nutrition Foundation Nutrition Bulletin* 30, pp. 27-54.

O'Dea K., Nestel, P. J., and Antonoff, L. (1980). Physical Factors Influencing Postprandial Glucose and Insulin Responses to Starch. *American Journal of Clinical Nutrition* 33(4), pp. 760-765.

Pokorny, J. (1999). Changes of nutrients at frying temperatures. *In Frying of Food. Oxidation, Nutrient and Non-nutrient Antioxidants, Biologically Active Compounds*

and High Temperatures. Lancaster, Pennsylvania, USA; Technomic Publishing Company, Inc., pp. 69-95.

Saguy, I. S. & Pinthus, I. J. (1995). Oil Uptake during Deep-fat Frying: Factors and Mechanism. *Food Technology* 49(4), pp. 142-145, 152.

Saguy, I. S. & Dana, D. (2003). Integrated Approach to Deep Fat Frying: Engineering, Nutrition, Health and Consumer Aspects. *Journal of Food Engineering* 56, pp. 143-152.

Sajilata, M. G., Singhal, R. S., and Kulkarni, P. R. (2006). Resistant Starch – A Review. *Comprehensive Reviews in Food Science and Food Safety* 5, pp. 1-17

Sanz, T., Salvador, A. & Fiszman, S. M. (2007). Performance of three different types of resistant starch in fried battered food. *European Food Research and Technology* 227(1), pp. 21-27.

Sanz, T., Salvador, A., & Fiszman, S. M. (2008). Resistant Starch (RS) in Battered Fried Products: Functionality and High-Fibre Benefit. *Food Hydrocolloids* 22(4), pp. 543-549.

Shih, F. and Daigle, K. 1999. Oil Uptake Properties of Fried Batters from Rice Flour. *Journal of Agricultural and Food Chemistry* 47(4), pp. 1611-1615.

Srichuwong, S. (2006). Starches from Different Plant Origins: From Structure to Physicochemical Properties. Japan: Mie University.

Vania, L. (2017). Application of Resistant Starch from Unripe "Kepok" Banana (*Musa paradisiaca formatypica*) in Batter Coating Formulation to Reduce Oil Absorption in Fried Food. BS Thesis. Department of Food Technology. Swiss German University, Tangerang, Indonesia.

Vatanasuchart, N.Niyomwit , B. Wongkrajang, K. (2012). Resistant starch content, in vitro starch digestibility and physico-chemical properties of flour and starch from Thai bananas. *Maejo Int. J. Sci. Technol.* 6(02), pp. 259-271.

Zhang, P., Whistler, R., BeMiller, J., & Hamaker, B. (2005). Banana starch: production, physicochemical properties, and digestibility—a review. *Carbohydrate Polymers* 59(4), pp. 443-458.



REFERENCES for Table 2.2

BSN. (2012). *Tempe: Persembahan Indonesia untuk Dunia* (SNI 3144:2009). Jakarta: Badan Standardisasi Nasional.

Maghribi, F. G. U. (2016). *Profil Asam Lemak, Kolesterol dan Jaringan Fillet Ikan Baronang (Siganus sp.) Akibat Proses Penggorengan*. Bogor: Institut Pertanian Bogor.

Mahmudatussa'adah A. (2014). *Chemical Composition of Cilembu Sweet Potato (Ipomoea batatas L) at Various Storage Time as Raw Material of Liquid Sugar*. Bandung: Universitas Pendidikan Indonesia.

Midayanto, D. N. & Yuwono, S. S. (2014). *Penentuan Atribut Mutu Tekstur Tahu untuk Direkomendasikan sebagai Syarat Tambahan dalam Standar Nasional Indonesia. Jurnal Pangan dan Agroindustri* 2(4), pp. 259-267.

Mitra, J., Shrivastava, S. L., & Rao, P. S. (2012). Onion Dehydration: A Review. *Journal of Food Science and Technology* 49(3), pp. 267-277.

Koswara, S. *Modul Teknologi Pengolahan Umbi-umbian Bagian 6: Pengolahan Singkong*. SEAFast Center Bogor Agricultural University.

Putri, A. R. (2012). *Pengaruh Kadar Air terhadap Tektur dan Warna Keripik Pisang Kepok (Musa parasidiaca formatypica)*. Makassar: Universitas Hasanuddin.

Rahardjo, S. (2003). *Kajian Proses dan Formulasi Pembuatan Sosis Nabati dari Jamur Tiram Putih (Pleurotus ostreatus)*. Bogor: Institut Pertanian Bogor.

Trilaksani, W., Erungan, A. C., & Mardi, S. (2004). *Pengaruh Suhu dan Lama Pengovenan terhadap Karakteristik Cumi-cumi (Loligo sp) Kertas. Buletin Teknologi Hasil Perikanan VIII (II)*, pp. 19-29.

Tropical Fruit Farm – “Cempedak Assorted (*Artocarpus integer*)”
<http://www.tropicalfruits.com.my/pdf/Cempedak-k.pdf/>, accessed on November 30, 2017.

Wijaya, M. G. (2015). *Karakteristik Kandungan Gizi Udang Vannamei (Litopenaeus vannamei) dari Sistem Budidaya yang Berbeda*. Bogor: Institut Pertanian Bogor.

Wala, J., Ransaleleh, T., Wahyuni, I., & Rotinsulu, M. (2016). Kadar air, pH dan Total Mikroba Daging Ayam yang Ditambahkan Kunyit Putih (*Curcuma mangga Val.*). *Jurnal ZooteK* 36(2), pp. 405-416.



SWISS GERMAN UNIVERSITY